

Aus der Klinik für Psychiatrie, Neurologie, Psychosomatik und Psychotherapie im  
Kindes- und Jugendalter der Universität Rostock

Direktor: Prof. Dr. med. F. Häßler

und aus der

Universitätsfrauenklinik und Poliklinik am Klinikum Südstadt Rostock

Direktor: Prof. Dr. med. B. Gerber

**Schwangerschaft und Geburt im minderjährigen Alter –  
Ist die Risikobelastung ein Effekt des Alters oder bedingt durch  
Bildungsstand, Partnerstatus und/oder Tabakkonsum?  
Eine Matched - Pairs - Studie an Erstgebärenden der  
Jahrgänge 2000 – 2004  
- Band 2: Anhang -**

---

Inauguraldissertation

zur

Erlangung des akademischen Grades

Doktor der Medizin

der Medizinischen Fakultät

der Universität Rostock

vorgelegt von

Sara Fracassi, geb. am 16.5.1981 in Bergisch Gladbach

aus Overath

verteidigt am 8.7.2009

Dekan: Prof. Dr. med. E. Reisinger

- Gutachter:
1. Prof. Dr. med. habil. Frank Häbler  
Klinik für Psychiatrie, Neurologie, Psychosomatik und  
Psychotherapie im Kindes- und Jugendalter der Universität  
Rostock
  2. Prof. Dr. med. habil. Bernd Gerber,  
Universitätsfrauenklinik und Poliklinik am  
Klinikum Südstadt Rostock
  3. Dr. med. Oliver Bilke,  
Klinik für Kinder-Jugendpsychiatrie und Psychotherapie,  
Vivantes Klinikum Hellersdorf

---

## **Inhaltsverzeichnis**

<b>Anhang.....</b>	<b>1</b>
A1. Risikoliste .....	1
A2. Vergleichslisten der Matched-Pairs 2000 – 2004.....	16
A3. Übersicht der Ergebnisse .....	25
A3.1 Anamnestische Angaben vor der Schwangerschaft.....	25
A3.2 Schwangerschaftsverlauf.....	41
A3.3 Geburt.....	67
A3.4 Kindliches Outcome .....	89
A3.5 Jugendamt oder Vormundschaft .....	115
A3.6 Diskussion.....	117



---

## Anhang

### A1. Risikoliste

Vom Untersucher einzutragen	
Datum Aktenstudium:	durchgeführt durch:
Interviewdatum:	durchgeführt durch:
Mimücode:	Code der Matchmutter:
Geburtsdatum Mutter:	Alter:
Geburtsdatum des Kindes:	

Notfall :

Konfession:

Akte Inter-  
wiew001 Familienstand:
 

- 01 ledig
- 02 geschieden
- 03 verheiratet
- 04 eheähnliche Gemeinschaft

002 Beruf vor der Schwangerschaft
 

- 01 ungelernter Arbeiter
- 02 angelernter Berufe
- 03 Facharbeiter, Handwerker, Angestellte, Beamte im Dienst
- 04 mittlere qualifizierte Angestellte, Beamte im einfachem Dienst
- 05 höher qualifizierte Angestellte, Beamte im gehobenen Dienst
- 06 leitende Angestellte, Beamte im höheren Dienst
- 07 Kleinste Selbständige, ambulantes Gewerbe
- 08 kleine selbständige Gewerbetreibende
- 09 selbständige Handwerker, Landwirte, Gewerbetreibende (kl. Betriebe)
- 10 selbständige Handwerker, Landwirte (mittl. Betriebe)
- 11 selbständige Akademiker, freie Berufe, (größere Unternehmen)
- 12 arbeitslos
- 13 Schülerin
- 14 Berufsvorbereitungsjahr
- 15 Azubi

003 Welchen Beruf üben Sie aus?

## Fragen zum Kindsvater

004 Welchen Beruf übt der Kindsvater derzeitig aus?

- 01 ungelernter Arbeiter
- 02 angelernter Berufe
- 03 Facharbeiter, Handwerker, Angestellte, Beamte im Dienst
- 04 mittlere qualifizierte Angestellte, Beamte im einfachem Dienst
- 05 höher qualifizierte Angestellte, Beamte im gehobenen Dienst
- 06 leitende Angestellte, Beamte im höheren Dienst
- 07 Kleinste Selbständige, ambulantes Gewerbe
- 08 kleine selbständige Gewerbetreibende
- 09 selbständige Handwerker, Landwirte, Gewerbetreibende (kl. Betriebe)
- 10 selbständige Handwerker, Landwirte (mittl. Betriebe)
- 11 selbständige Akademiker, freie Berufe, (größere Unternehmen)
- 12 arbeitslos
- 13 Schüler
- 14 Berufsvorbereitungsjahr
- 15 Azubi

## Risiken - Kindesmutter

005 Rauchen Sie?

006 Wenn ja  
Wie viele Zigaretten rauchen Sie pro Tag?

- 01 weniger als 10
- 02 10-30
- 03 mehr als 30

007 Haben Sie während der Schwangerschaft geraucht?

007 Wenn ja  
Wie viele Zigaretten pro Tag?

- 01 Weniger als 10
- 02. 20-30
- 03 mehr als 30

---

009 Leiden Sie an einer dieser Erkrankungen?

- 01 Herz-Kreislauf-Erkrankungen
- 02 chronische Gelenkerkrankungen
- 03 chronische Lungenerkrankungen
- 04 chronische Magendarmerkrankungen
- 05 Geschlechtskrankheiten
- 06 Stoffwechselerkrankungen
- 06.1 Diabetes
- 07 chronische Schmerzen
- 08 Migräne
- 09 Gynäkologische Erkrankungen
- 10 Leber oder Galle Erkrankungen
- 11 andere, welche: \_\_\_\_\_
- 12 Allergien
- 13 Tumore \_\_\_\_\_

010 Familienanamnestische Erkrankungen

- 01 Herz-Kreislauf-Erkrankungen
- 02 chronische Gelenkerkrankungen
- 03 chronische Lungenerkrankungen
- 04 chronische Magendarmerkrankungen
- 05 Geschlechtskrankheiten
- 06 Stoffwechselerkrankungen
- 06.1 Diabetes
- 07 chronische Schmerzen
- 08 Migräne
- 09 Gynäkologische Erkrankungen
- 10 Leber oder Galle Erkrankungen
- 11 andere, welche: \_\_\_\_\_
- 12 Allergien
- 13 Tumore \_\_\_\_\_

011 Wurden Sie vor der Schwangerschaft gynäkologisch operiert?

012 Wenn ja  
Sind Sie in regelmäßiger Behandlung?

- 013 Waren eine dieser psychischen Erkrankungen vor der Schwangerschaft bei Ihnen bekannt?
- 01 F1 Sucht
  - 02 F2 Psychosen
  - 03 F3 Depression
  - 04 F4 Anpassungsstörungen
  - 05 F5 Anorexie  
Bulimi
  - 06 F6 Persönlichkeitsstörungen
  - 07 F7 Intellektueller Leistungsabfall
  - 08 F8 Teilleistungsstörungen/ Entwicklungsstörungen
  - 09 F9 Störungen des Sozialverhaltens / Hyperkinetische Störungen
- 014 Wurde die Erkrankung diagnostiziert?
- 015 Wenn ja  
Wie wurde die Erkrankung behandelt?
- 01 Wurde nicht behandelt
  - 02 Psychotherapeutische Therapie
  - 03 Medikamentöse Therapie
- 016 Haben sie vor der Schwangerschaft
- 01 Sich selbst verletzt
  - 02 Drogen konsumiert
- 017 Im Folgenden frage ich Sie nach dem gegenwärtigen Konsum verschiedener Drogen.



**in den letzten 4 Wochen**

	jemals konsumiert	Einstiegsalter	gelegentlich 1-3 mal	Wochenendkonsum 4-8 mal	regelmäßig Konsum je Woche 3-6 mal	täglicher Konsum	längste Zeit des Nichtgebrauchs		
							Wochen	Monate	Jahre
Tabak									
Alkohol									
Cannabinoide									
Pilze									
Gase									
Schnüffelstoffe									
Stimulanzien (Speed)									
Entaktogene (Ecstasy)									
Kokain									
Crack									
Halluzinogene: LSD									
Opioide									
Methadon									
Medikamente									

018 Nehmen Sie zur Zeit ärztlich verordnete Medikamente ein?

019 Ist nach der Geburt eine Erkrankungen aufgetreten?

- 01 ja  
02 nein  
03 was

**Schwangerschaftsanamnese**

020 Gewicht zu Schwangerschaftsbeginn

021 Körpergröße:

022 Gewichtszunahme in der gesamten Schwangerschaft

023 Anzahl der Abruption?

024 Gab es Komplikationen bei oder nach der Abruption?

- 01 Übelkeit

- 02 allergische Reaktion
- 03 Atemnot und Brustschmerz
- 04 pathologische Nachblutungen
- 05 Infektion der Gebärmutter
- 06 Durchstoßung der Gebärmutter
- 06 Sterilität nach Verklebung der Eileiter aufgrund einer Infektion

025 Anzahl der Fehlgeburten?

026 Anti D Prophylaxe?

### **Schwangerschaftsverlauf**

027 Anzahl der Ultraschalluntersuchungen

028 Infektionskrankheiten

- 01 Harnwegsinfekte
- 02 Röteln
- 03 CMV
- 04 Toxoplasmose
- 05 Herpes genitalis Typ 2
- 06 HIV
- 07 Treponema pallidum
- 08 Chlamydien
- 09 HPV
- 10 Hepatitis B
- 11 Hepatitis C
- 12 Pilze
- 13 B- Streptokokken

029 Andere Infektionskrankheiten

- 01 der Atemwege
- 02 Nierenbeckenentzündung
- 03 Magendarmerkrankung
- 04 Varizellen
- 05 Parvovirus
- 06 andere: \_\_\_\_\_

030 Isolierte Hypertonie

031 Präeklampsie

032 Eklampsie

033 HELLP-Syndrom

- 
- 034 Hypotensives Syndrom
- 035 Amnioninfusionssyndrom (Fruchtwasserembolie)
- 036 Amnioninfektionsyndrom (Eihäuteentzündung)
- 037 Haben Sie während der SS an einer dieser Erkrankungen gelitten?
- 01 Plazentainsuffizienz  
02 Zervixinsuffizienz  
03 Emesis  
04 Andere
- 038 Blutungen
- 01 Erste Schwangerschaftshälfte  
02 Zweite Schwangerschaftshälfte  
03 Pathologische Blutung unter Geburt
- 039 Ist während der Schwangerschaft eine dieser Erkrankungen aufgetreten?
- 01 Herzkreislauferkrankung  
02 chronische Gelenkerkrankungen  
03 chronische Lungenerkrankungen  
04 chronische Magendarmerkrankungen  
05 Geschlechtskrankheiten  
06 Stoffwechselerkrankungen  
07 chronische Schmerzen  
08 Migräne  
09 Gynäkologische Erkrankungen  
10 Leber oder Galle Erkrankungen  
11 andere: \_\_\_\_\_  
12 Allergien  
13 Tumore \_\_\_\_\_
- 040 Waren Sie während der Schwangerschaft im Krankenhaus?
- 041 Wenn ja  
Warum?
- 042 Wenn ja.  
Wie lange? (Angabe in Wochen)
- 043 Nahmen sie im Krankenhaus Medikamente ein?
-

- 
- 01 Spasmolytika
  - 02 Analgetika
  - 03 Allgemeinanalgesie
  - 04 Lokalanästhesie
  - 05 Regionalanästhesie
  - 06 Oxytocin
  - 07 Tokolytika
  - 08 Antibiotika
  - 09 anderes( Prophylaxe)

- 044 Ikterus während der Schwangerschaft?
- 045 Wenn ja  
Wie hoch war der Bilirubinwert?
- 046 Gab es chirurgische Eingriffe während der Schwangerschaft?
- 047 Wenn ja.  
Welcher Eingriff wurde vorgenommen?
- 048 Hatten Sie Unfälle während der Schwangerschaft?
- 049 Wenn ja.  
Welche ?

### Serologische Tests & Frühdiagnostik

- 050 Amniozentese
- 051 Wenn ja.  
Gründe für die Amniozentese?
- 052 Chorionzottenbiopsie
- 053 Wenn ja.  
Warum?
- 054 Wenn Komplikationen:  
Welche?
- 055 Wenn ja  
Warum?
- 056 Wenn ja

---

Anomalien des Kindes

- 01 Gliedmaßen
- 02 Innere Organe
- 03 Schädel
- 04 Fruchtwasser
- 05 Magen? Darm
- 06 Zwerchfell

**Geburt**057 In welcher Woche haben Sie entbunden?058 Art der Entbindung

- 01 Spontan
- 02 Sectio
- 03 Vakuumextraktion
- 05 Forceps

059 Lage

- 01 Beckenlage
- 02 Schädellage
- 03 Querlage

060 Wenn sectio aus welchen Grund?

- 01 primär
- 02 sekundär

061 Wenn sekundär

- 01 pathologisches CTG
- 02 fetopelvines Missverhältnis
- 03 protrahierte EP
- 04 andere: \_\_\_\_\_

062 Beginn der Geburt

- 01 spontan
- 02 wehenfördernde Mittel
- 03 priming

- 
- 063 Wann war der Blasensprung
- 01 Unter 4h vor Geburt  
02 zwischen 4-12h vor Geburt  
03 Über 12h vor Geburt  
04 unter der Geburt
- 064 Blasensprung
- 01 spontan  
02 künstlich
- 065 Fruchtwasser
- 01 klar  
02 grün  
03 blutig
- 066 Dauer der Geburt
- 067 Kindseitige Komplikationen
- 01 abnorme fetale HF  
02 Nabelschnurumschlingung  
03 Insertio der Nabelschnur  
04 Kephalhämatom  
05 Caput succedaneum  
06 Obere Plexuslähmung  
07 Untere Plexuslähmung  
08 Klavikulafraktur  
09 Konjunktivitis  
10 andere \_\_\_\_\_
- 068 Medikamentengabe unter Geburt
- 01 Spasmolytika  
02 Analgetika  
03 Allgemeinanalgesie  
04 Lokalanästhesie  
05 Regionalanästhesie  
06 Oxytocin  
07 Tokolytika  
08 Antibiotika  
09 anderes u.a.Prophylaxe
- 069 War die Geburt Ihres Kindes mit Komplikationen verbunden?
-

- 
- 070 Wenn ja  
Welche Komplikationen?
- 01 Plazenta praevia
  - 02 tiefer Sitz der Plazenta
  - 03 vorzeitige Plazentaablösung
  - 04 Randsinusblutung
  - 05 Uterusruptur
  - 06 Gefäßruptur bei Insertio velamentosa
  - 07 Ruptur eines Varixknotens der Scheide
  - 08 Dammriss/ Dammschnitt
  - 09 Scheidenriss
  - 10 Zervixriss
  - 11 Schock( kardiogener, posthämorrhagischer, vasomotorischer)
  - 12 Atonie
  - 13 andere \_\_\_\_\_
- 071 Blutverlust
- 01 0-250ml
  - 02 250-500ml
  - 03 mehr als 500ml
- 072 Verweildauer auf der Entbindungsstation
- 01 unter 3 Tage
  - 02 3-5 Tage
  - 03 5-10 Tage
  - 04 über 10 Tage
  - 05 Entlassung gegen ärztlichen Rat
- 073 Gab es Komplikationen im Wochenbett?
- 074 Wenn ja  
Welche Komplikationen?
- 01 Toxisches Schocksyndrom
  - 02 Fieber ( Puerperale Infektion)
  - 03 Wundheilungsstörungen
  - 04 Mastitis
  - 05 Wochenbettdepression
  - 06 Thrombose
  - 07 Embolie
  - 08 Anämie

---

075 Von der Norm abweichende Laborwerte bei Kindesmutter?

01 Leberwerte  
 02 Eisenwerte  
 03 Gerinnungswerte  
 04 Cholesterin, LDL, HDL  
 05 Triglyceride  
 06 Bilirubin  
 07 andere \_\_\_\_\_

076 Haben sie gestillt?

077 Wenn nicht aus welchen Grund?

01 Wunsch  
 02 ärztliche Indikation

### Neugeborenenangaben

078 Geschlecht:

01 Mädchen  
 02 Junge

079 Größe

080 Gewicht

081 Kopfumfang

082 APGAR nach 1 min

083 APGAR nach 5 min

084 APGAR nach 10 min

085 Sauerstoffgabe

086 Medikamentöse Soforttherapie?

087 Wenn ja  
Welche medikamentöse Soforttherapie?

088 Wurden Reanimationsmaßnahmen eingeleitet?



- 
- 089 Reifestatus des Neugeborenen
- 01 eutroph  
02 hypertroph  
03 hypotroph
- 090 Reifestatus des Frühgeborenen
- 01 eutroph  
02 hypertroph  
03 hypotroph
- 091 Unphysiologischer Ikterus neonatorum
- 01 über 12,5mg\dl  
02 über 15mg\ dl
- 092 Lungenreifung erfolgt?
- 093 Wenn ja,   
Wer?
- 01 Mutter  
02 Kind
- 094 Verweildauer auf Station
- 01 unter 3 Tage  
02 3-5 Tage  
03 5-10 tage  
04 über 10 Tage  
05 Entlassen gegen ärztlichen Rat
- 095 War das Neugeborene auf der neonatologischen Station?
- 096 Wenn ja wie lange?
- 01 unter 3 Tage  
02 3-5 Tage  
03 5-10 tage  
04 über 10 Tage
- 097 Wenn ja   
Wurde eine Polysomnographie durchgeführt?
-

---

098	<u>PH-Wert des Kindes</u>	<input type="checkbox"/> <input type="checkbox"/>
	01 > 7,3	
	02 <7,2 bis 7,3	
	03 7,1 bis 7,2	
	04 unter 7,1	
099	<u>Base Excess (BE)</u>	<input type="checkbox"/> <input type="checkbox"/>
	01 +5-(-5)	
	02 -5-(-10)	
	03 -10-(-15)	
	04 -15-(-20)	
	05 -20-(-30)	
	06 mehr als 30	
100	<u>Kopfhaare abgegrenzt?</u>	<input type="checkbox"/> <input type="checkbox"/>
101	<u>Lanugobehaarung?</u>	<input type="checkbox"/> <input type="checkbox"/>
102	<u>Fingernägel überragend</u>	<input type="checkbox"/> <input type="checkbox"/>
103	<u>Testes eingetreten</u>	<input type="checkbox"/> <input type="checkbox"/>
104	<u>Labien geschlossen</u>	<input type="checkbox"/> <input type="checkbox"/>
105	<u>Überreifezeichen</u>	<input type="checkbox"/> <input type="checkbox"/>
106	<u>Ergebnis der U2</u>	<input type="checkbox"/> <input type="checkbox"/>
<hr/>		
107	<u>Ergebnis der Hüftsonographie (Einteilung nach Graf)</u>	<input type="checkbox"/> <input type="checkbox"/>
	01 oB	
	02 kontrollbedürftig	
	03 pathologisch	
108	Wenn 03 <u>Welche Störung war vorhanden?</u>	<input type="checkbox"/> <input type="checkbox"/>
	01 angeborene Hüftluxation	
	02 Hüftsubluxation	
	03 Hüftdysplasie	

---

- 
- |     |   |  |
|-----|---|--|
| 109 | <u>Ergebnis Schädelsonographie</u>                        | <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> |
|     | 01 in Ordnung   |  |
|     | 02 pathologisch   |  |
| 110 | <u>Welche Pathologie fiel auf?</u>                        | <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> |
| 111 | <u>Von der Norm abweichende Laborparameter beim Kind?</u> | <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> |
|     | 01 Leberwerte   |  |
|     | 02 Eisenwerte   |  |
|     | 03 Gerinnungswerte  |  |
|     | 04 Cholesterin, LDL, HDL, Triglyceride                    |  |
|     | 05 Bilirubin  |  |
|     | 06 Glucose  |  |
|     | 07 Harnstoff  |  |
|     | 08 andere   |  |
| 112 | <u>Ergebnis Hörscreening</u>                              | <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> |
|     | 01 in Ordnung   |  |
|     | 02 pathologisch   |  |
| 113 | <u>Neugeborenenenscreening</u>                            | <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> |
|     | 01 in Ordnung   |  |
|     | 02 steht aus  |  |
|     | 03 pathologisch hinsichtlich.....                         |  |
| 114 | <u>Weitere pathologische Befunde?</u>                     | <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> |

Codierung: 0 – Nein; 01 – Ja ; 88 – trifft nicht zu; 99 - unbekannt

## A2. Vergleichslisten der Matched-Pairs 2000 – 2004

		2000		
Müttercode	Erstgebärend	Partnerstatus	Schulbildung	Rauchen
1) MV 85	*	ledig	Schülerin	< 10
KN 82	*	ledig	arbeitslos	< 10
2) CL 82	*	ledig	Schülerin	<10
GM 81	*	ledig	arbeitslos	< 10
3) DW 82	*	ledig (KV)	Schülerin	Nein
GM 77	*	ledig (KV)	arbeitslos	Nein
4) NS 82	*	ledig (KV)	Schülerin	< 10
HS 74	*	ledig (KV)	arbeitslos	< 10
5) SB 85	*	ledig (KV)	Schülerin	10 – 30
HP 80	*	ledig (KV)	arbeitslos	10 – 30
6) MV 84	*	ledig	Schülerin	< 10
KA 78	*	ledig	arbeitslos	<10
7) NZ 82	*	ledig	Hausfrau	Nein
PB 79	*	ledig	arbeitslos	Nein
8) YT 83	*	ledig (KV)	Schülerin	Nein
SA 80	*	ledig (KV)	arbeitslos	Nein
9) EK 84	*	ledig	Schülerin	Nein
DML 70	*	ledig	arbeitslos	Nein
10) DW 82	*	ledig (KV)	Schülerin	Nein
LN 80	*	ledig (KV)	arbeitslos	Nein
11) EK 83	*	ledig	Schülerin	Nein
ST 79	*	ledig	arbeitslos	Nein
12) MR 83	*	ledig (KV)	arbeitslos	< 10
LM 79	*	ledig (KV)	berufsunfähig	< 10
13) RT 83	*	ledig	AZUBI	10 – 30
LD 81	*	ledig	arbeitslos	10 – 30
14) SB 83	*	ledig (KV)	Schülerin	< 10
PA 80	*	ledig (KV)	arbeitslos	< 10
15) BL 82	*	?	arbeitslos	< 10
RY 77	*	ledig (KV)	arbeitslos	< 10

---

Müttercode	Erstgebärend	2000		
		Partnerstatus	Schulbildung	Rauchen
16) VE 84	*	ledig (KV)	?	Nein
RN 78	*	ledig (KV)	?	Nein
17) NS 82	*	ledig	arbeitslos	10 – 30
SC 78	*	ledig	arbeitslos	10 – 30
18) EWO 83	*	ledig	Schülerin	Nein
BA 79	*	ledig	arbeitslos	Nein
19) CL 82	*	ledig (KV)	arbeitslos	10 – 30
PS 75	*	ledig (KV)	arbeitslos	10 – 30
20) KK 83	*	ledig	arbeitslos	Nein
WF 81	*	ledig	arbeitslos	Nein
21) CH 85	*	ledig	Schülerin	< 10
LN 80	*	ledig	arbeitslos	< 10
22) CS 83	*	ledig	AZUBI	10 – 30
JB 79	*	ledig	arbeitslos	10 – 30

Sofern die Akten Daten zum Zusammenleben der jeweiligen Mütter mit dem Kindsvater lieferten, wurde die in der Tabelle ledig (KV) gekennzeichnet.

		2001		
Müttercode	Erstgebärend	Partnerstatus	Schulbildung	Rauchen
1) SS 84	*	ledig	AZUBI	Nein
FA 83	*	ledig	Praktikantin	Nein
2) LK 83	*	ledig	Schülerin	Nein
RD 81	*	ledig	arbeitslos	Nein
3) SR 83	*	ledig (KV)	arbeitslos	< 10
ED 78	*	ledig (KV)	arbeitslos	< 10
4) JU 83	*	ledig (KV)	AZUBI	Nein
LR 79	*	ledig (KV)	AZUBI	Nein
5) CK 84	*	ledig	Schülerin	Nein
TA 81	*	ledig	arbeitslos	Nein
6) SL 83	*	ledig	Schülerin	< 10
BJ 81	*	ledig	ABM	<10
7) SL 83	*	ledig	arbeitslos	< 10
BC 83	*	ledig	arbeitslos	< 10
8) DT 83	*	ledig	Schülerin	Nein
SS 76	*	ledig	Hausfrau	Nein
9) CR 83	*	ledig	AZUBI	< 10
FA 82	*	ledig	AZUBI	< 10
10) SE 84	*	ledig	arbeitslos	< 10
SR 67	*	ledig	arbeitslos	< 10
11) MS 83	*	ledig	arbeitslos	< 10
CM 81	*	ledig	arbeitslos	< 10
12) UB 86	*	ledig	Schülerin	Nein
BY 75	*	ledig	arbeitslos	Nein
13) MW 84	*	ledig	Schülerin	Nein
AD 80	*	ledig	arbeitslos	Nein
14) EN 84	*	ledig	Schülerin	Nein
SB 77	*	ledig	arbeitslos	Nein
15) SP 83	*	ledig (KV)	AZUBI	< 10
BA 81	*	ledig (KV)	AZUBI	< 10
16) SD 85	*	ledig (KV)	Schülerin	Nein
KD 79	*	ledig (KV)	arbeitslos	Nein
17) DS 83	*	ledig	Schülerin	< 10
HY 83	*	ledig	arbeitslos	< 10
18) JB 83	*	ledig (KV)	AZUBI	< 10
SM 80	*	ledig (KV)	AZUBI	< 10

---

		2001		
Müttercode	Erstgebärend	Partnerstatus	Schulbildung	Rauchen
19) UK 84	*	ledig	arbeitslos	Nein
RN 78	*	ledig	arbeitslos	Nein
20) PS 85	*	ledig (KV)	arbeitslos	10 – 30
AU 72	*	ledig (KV)	arbeitslos	10 – 30
21) HJ 84	*	ledig	Hausfrau	< 10
FN 78	*	ledig	arbeitslos	< 10
22) KB 85	*	ledig	Schülerin	Nein
US 81	*	ledig	AZUBI	Nein
23) CG 84	*	ledig	?	< 10
KN 81	*	ledig	AZUBI	< 10
24) DB 84	*	ledig	?	10 – 30
TN 83	*	ledig	Schülerin	10 – 30

		2002		
Müttercode	Erstgebärend	Partnerstatus	Schulbildung	Rauchen
1) NG 84	*	ledig	?	< 10
SK 82	*	ledig	Verkäuferin	< 10
2) MW 84	*	ledig (KV)	AZUBI	10 – 30
WS 74	*	ledig (KV)	Friseurin	10 – 30
3) AK 85	*	ledig	Schülerin	Nein
TE 80	*	ledig	Hausfrau	Nein
4) KW 84	*	ledig	arbeitslos	10 – 30
OJ 83	*	ledig	arbeitslos	10 – 30
5) MR 85	*	ledig (KV)	AZUBI	Nein
HD 83	*	ledig (KV)	arbeitslos	Nein
6) SS 85	*	ledig	arbeitslos	10 – 30
QJ 71	*	ledig	arbeitslos	10 – 30
7) CA 85	*	ledig	AZUBI	< 10
LI 81	*	ledig	AZUBI	< 10
8) JK 85	*	ledig	BVJ	Nein
HR 83	*	ledig	BVJ	Nein
9) NS 85	*	ledig	Schülerin	< 10
HN 79	*	ledig	arbeitslos	< 10
10) AKK 86	*	ledig (KV)	Schülerin	< 10
LC 80	*	ledig (KV)	arbeitslos	< 10
11) SN 85	*	ledig (KV)	Schülerin	< 10
RN 77	*	ledig (KV)	arbeitslos	< 10
12) SF 86	*	ledig (KV)	?	10 – 30
RS 83	*	ledig (KV)	arbeitslos	10 – 30
13) JG 88	*	ledig	Schülerin	< 10
HJ 82	*	ledig	arbeitslos	< 10
14) SW 85	*	ledig	arbeitslos	Nein
FN 80	*	ledig	Hausfrau	Nein
15) NN 85	*	ledig	AZUBI	10 – 30
SN 82	*	ledig	AZUBI	10 – 30
16) MP 85	*	ledig (KV)	Verkäuferin	< 10
GC 78	*	ledig (KV)	Verkäuferin	< 10



---

		2002		
Müttercode	Erstgebärend	Partnerstatus	Schulbildung	Rauchen
17) MK 85	*	ledig	AZUBI	10 – 30
HY 83	*	ledig	arbeitslos	10 – 30
18) SS 85	*	ledig	Schülerin	Nein
SC 83	*	ledig	arbeitslos	Nein
19) KS 84	*	ledig (KV)	AZUBI	Nein
BE 82	*	ledig (KV)	AZUBI	Nein
20) DJ 86	*	ledig (KV)	Schülerin	10 – 30
WS 74	*	ledig (KV)	arbeitslos	10 – 30

		2003		
Müttercode	Erstgebärend	Partnerstatus	Schulbildung	Rauchen
1) NK 86	*	ledig	AZUBI	Nein
BD 83	*	ledig	AZUBI	Nein
2) AK 85	*	ledig	Schülerin	< 10
KU 82	*	ledig	arbeitslos	< 10
3) RK 85	*	ledig	AZUBI	< 10
SM 82	*	ledig	AZUBI	< 10
4) OS 85	*	ledig (KV)	Schülerin	Nein
SV 85	*	ledig (KV)	Schülerin	Nein
5) NM 86	*	ledig	Schülerin	< 10
LY 78	*	ledig	arbeitslos	< 10
6) AR 85	*	ledig	arbeitslos	< 10
KC 83	*	ledig	arbeitslos	< 10
7) WS 85	*	ledig	AZUBI	< 10
RD 81	*	ledig	AZUBI	< 10
8) NK 87	*	ledig	Schülerin	Nein
BE 80	*	ledig	arbeitslos	Nein
9) NL 85	*	ledig	AZUBI	Nein
KK85	*	ledig	AZUBI	Nein
10) NF 88	*	ledig (KV)	Schülerin	< 10
MK 83	*	ledig (KV)	arbeitslos	< 10
11) AR 86	*	verheiratet	arbeitslos	Nein
FB 76	*	verheiratet	arbeitslos	Nein
12) AH 86	*	ledig (KV)	BVJ	10 – 30
SS 82	*	ledig (KV)	AZUBI	10 – 30
13) CB 87	*	ledig (KV)	arbeitslos	Nein
SK 79	*	ledig (KV)	arbeitslos	Nein
14) NB 87	*	ledig	Schülerin	10 – 30
WS 84	*	ledig	AZUBI	10 – 30
15) AB 88	*	ledig	Schülerin	Nein
KJ 77	*	ledig	arbeitslos	Nein
16) MS 87	*	?	Schülerin	Nein
SA 79	*	ledig	arbeitslos	Nein

---

Müttercode	Erstgebärend	2003		
		Partnerstatus	Schulbildung	Rauchen
17) FH 87	*	ledig	Schülerin	Nein
FN 82	*	ledig	arbeitslos	Nein
18) SS 85	*	ledig	AZUBI	< 10
SC 83	*	ledig	Hausfrau	< 10
19) PS 86	*	verheiratet	arbeitslos	Nein
PK 76	*	verheiratet	arbeitslos	Nein
20) AT 88	*	ledig	Schülerin	?
BJ 82	*	ledig	arbeitslos	Nein
21) MP 86	*	ledig	?	Nein
TC 77	*	ledig	?	Nein
22) NH 86	*	ledig	arbeitslos	< 10
HB 82	*	ledig	arbeitslos	< 10
23) AW 86	*	ledig	Schülerin	< 10
BJ 81	*	ledig	arbeitslos	< 10
24) BM 85	*	ledig	Schülerin	< 10
RC 80	*	ledig	arbeitslos	< 10

		2004		
Müttercode	Erstgebärend	Partnerstatus	Schulbildung	Rauchen
1) AM 89	*	?	Schülerin	Nein
KL 84	*	?	arbeitslos	Nein
2) AK 85	*	ledig	BVJ	< 10
SK 83	*	ledig	FSJ	< 10
3) NW 88	*	ledig	arbeitslos	< 10
BP 82	*	ledig	arbeitslos	< 10
4) CL 87	*	ledig (KV)	AZUBI	Nein
TW 85	*	ledig (KV)	AZUBI	Nein
5) AP 87	*	ledig	?	Nein
SL 83	*	ledig	?	Nein
6) HK 90	*	ledig	?	< 10
SC 85	*	ledig	?	< 10
7) MW 87	*	ledig	BVJ	< 10
SH 83	*	ledig	arbeitslos	< 10
8) DE 87	*	ledig	arbeitslos	Nein
AH 79	*	ledig	arbeitslos	Nein
9) CT 88	*	ledig	Schülerin	< 10
KK85	*	ledig	arbeitslos	< 10
10) AN 87	*	ledig	arbeitslos	< 10
MP 80	*	ledig	arbeitslos	< 10
11) JP 89	*	ledig	Schülerin	< 10
BM 83	*	ledig	arbeitslos	< 10
12) MN 86	*	ledig	arbeitslos	Nein
NC 74	*	ledig	arbeitslos	Nein
13) SK 87	*	ledig	Schülerin	Nein
FA 81	*	ledig	arbeitslos	Nein
14) KG 88	*	ledig	Schülerin	< 10
DT 78	*	ledig	arbeitslos	< 10
15) JF 86	*	ledig	AZUBI	Nein
CH 78	*	ledig	AZUBI	Nein
16) JS 86	*	ledig	Kosmetikerin	Nein
SA 78	*	ledig	PTA	Nein
17) NF 88	*	ledig	Schülerin	10 – 30
TS 82	*	ledig	arbeitslos	10 – 30
18) IK 87	*	ledig	Schülerin	Nein
AP 73	*	ledig	arbeitslos	Nein
19) CM 88	*	ledig	Schülerin	< 10
BS 81	*	ledig	arbeitslos	< 10

### A3. Übersicht der Ergebnisse

#### A3.1 Anamnestische Angaben vor der Schwangerschaft

##### Somatische Erkrankungen vor der Schwangerschaft

Gesamtanzahl der Erkrankungen vor der Schwangerschaft

	code	N	Mean	Std. Deviation	Std. Error Mean
Gesamtanzahl der Erkrankung vor Schwangerschaft	Mimü	110	,84	,904	,086
	Vomü	110	1,45	1,282	,122

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Gesamtanzahl der Erkrankung vor Schwangerschaft	Equal variances assumed	14,065	,000	-4,072	218	,000	-,609	,150	-,904	-,314
	Equal variances not assumed			-4,072	195,888	,000	-,609	,150	-,904	-,314

Gesamtanzahl der erkrankten Frauen

Crosstab

			Erkrankung		Total
			Nein	Ja	
code	Mimü	Count	61	49	110
		Expected Count	52,0	58,0	110,0
		% within code	55,5%	44,5%	100,0%
		% within Erkrankung	58,7%	42,2%	50,0%
	vomü	Count	43	67	110
		Expected Count	52,0	58,0	110,0
		% within code	39,1%	60,9%	100,0%
		% within Erkrankung	41,3%	57,8%	50,0%
Total		Count	104	116	220
		Expected Count	104,0	116,0	220,0
		% within code	47,3%	52,7%	100,0%
		% within Erkrankung	100,0%	100,0%	100,0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	5,908 <sup>a</sup>	1	,015	,021	,011	
Continuity Correction <sup>b</sup>	5,270	1	,022			
Likelihood Ratio	5,936	1	,015	,021	,011	
Fisher's Exact Test				,021	,011	
Linear-by-Linear Association	5,882 <sup>c</sup>	1	,015	,021	,011	,006
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 52,00.

c. The standardized statistic is 2,425.

## Herz-Kreislauf-Erkrankungen

## Count

		Herz Kreislauf Erkrankungen		Total
		Nein	Ja	
code	Mimü	105	5	110
	Vomü	107	3	110
Total		212	8	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,519 <sup>a</sup>	1	,471	,721	,361	
Continuity Correction <sup>b</sup>	,130	1	,719			
Likelihood Ratio	,524	1	,469	,721	,361	
Fisher's Exact Test				,721	,361	
Linear-by-Linear Association	,517 <sup>c</sup>	1	,472	,721	,361	,221
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 4,00.

c. The standardized statistic is -,719.

## chronische Gelenkerkrankungen

Count

		Chronische Gelenkerkrankungen		Total
		Nein	Ja	
code	Mimü	107	3	110
	Vomü	109	1	110
Total		216	4	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,019 <sup>a</sup>	1	,313	,622	,311	
Continuity Correction <sup>b</sup>	,255	1	,614			
Likelihood Ratio	1,065	1	,302	,622	,311	
Fisher's Exact Test				,372	,311	
Linear-by-Linear Association	1,014 <sup>c</sup>	1	,314	,622	,311	,250
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,00.

c. The standardized statistic is -1,007.

## chronische Lungenerkrankungen

Count

		Chronische Lungenerkrankungen		Total
		Nein	Ja	
code	Mimü	105	5	110
	Vomü	104	6	110
Total		209	11	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,096 <sup>a</sup>	1	,757	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,096	1	,757	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	,095 <sup>c</sup>	1	,758	1,000	,500	,231
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 5,50.

c. The standardized statistic is ,309.

## chronische Magen-Darmerkrankungen

Count

		Chronische Magen- Darmerkrankungen		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	109	1	110
Total		219	1	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is 1,000.

## Geschlechtskrankheiten



Count

---

		Geschlechts krankheiten	Total
		Nein	
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## Stoffwechselerkrankungen

Count

		Stoffwechselerkrankungen		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	108	2	110
Total		218	2	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,018 <sup>a</sup>	1	,155	,498	,249	
Continuity Correction <sup>b</sup>	,505	1	,477			
Likelihood Ratio	2,791	1	,095	,498	,249	
Fisher's Exact Test				,498	,249	
Linear-by-Linear Association	2,009 <sup>c</sup>	1	,156	,498	,249	,249
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is 1,417.

## Diabetes

Count

		Diabetes	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## chronische Schmerzen

Count

		chronische Schmerzen	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## Migräne

Count

		Migräne		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	106	4	110
Total		216	4	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,074 <sup>d</sup>	1	,044	,122	,061	
Continuity Correction <sup>e</sup>	2,292	1	,130			
Likelihood Ratio	5,619	1	,018	,122	,061	
Fisher's Exact Test				,122	,061	
Linear-by-Linear Association	4,056 <sup>c</sup>	1	,044	,122	,061	,061
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,00.

c. The standardized statistic is 2,014.

## Gynäkologische Erkrankungen

Count

		Gynäkologische Erkrankungen		Total
		Nein	Ja	
code	Mimü	109	1	110
	Vomü	104	6	110
Total		213	7	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,689 <sup>d</sup>	1	,055	,119	,060	
Continuity Correction <sup>e</sup>	2,361	1	,124			
Likelihood Ratio	4,080	1	,043	,119	,060	
Fisher's Exact Test				,119	,060	
Linear-by-Linear Association	3,672 <sup>c</sup>	1	,055	,119	,060	,052
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 3,50.

c. The standardized statistic is 1,916.

---

 Leber-/ Gallen- Erkrankungen

Count

		Leber oder Gallenerkrankungen		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	109	1	110
Total		218	1	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,995 <sup>a</sup>	1	,318	1,000	,502	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,382	1	,240	1,000	,502	
Fisher's Exact Test				1,000	,502	
Linear-by-Linear Association	,991 <sup>c</sup>	1	,320	1,000	,502	,502
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is ,995.

## Allergien

Count

		Allergien		Total
		Nein	Ja	
code	Mimü	80	30	110
	Vomü	79	31	110
Total		159	61	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,023 <sup>a</sup>	1	,880	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,023	1	,880	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	,023 <sup>c</sup>	1	,881	1,000	,500	,118
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 30,50.

c. The standardized statistic is ,150.

## Tumore

Count

		Tumore		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	107	3	110
Total		217	3	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,041 <sup>d</sup>	1	,081	,247	,123	
Continuity Correction <sup>e</sup>	1,352	1	,245			
Likelihood Ratio	4,200	1	,040	,247	,123	
Fisher's Exact Test				,247	,123	
Linear-by-Linear Association	3,028 <sup>c</sup>	1	,082	,247	,123	,123
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,50.

c. The standardized statistic is 1,740.

## andere Erkrankungen

Count

		andere Erkrankungen		Total
		Nein	Ja	
code	Mimü	95	15	110
	Vomü	63	47	110
Total		158	62	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	22,997 <sup>d</sup>	1	,000	,000	,000	
Continuity Correction <sup>e</sup>	21,582	1	,000			
Likelihood Ratio	23,867	1	,000	,000	,000	
Fisher's Exact Test				,000	,000	
Linear-by-Linear Association	22,893 <sup>c</sup>	1	,000	,000	,000	,000
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 31,00.

c. The standardized statistic is 4,785.

## Gynäkologische Operationen

Crosstab

			Gyn OP		Total
			Nein	Ja	
code	Mimü	Count	90	20	110
		Expected Count	91,5	18,5	110,0
		% within code	81,8%	18,2%	100,0%
		% within Gyn OP	49,2%	54,1%	50,0%
	vomü	Count	93	17	110
		Expected Count	91,5	18,5	110,0
		% within code	84,5%	15,5%	100,0%
		% within Gyn OP	50,8%	45,9%	50,0%
Total	Count	183	37	220	
	Expected Count	183,0	37,0	220,0	
	% within code	83,2%	16,8%	100,0%	
	% within Gyn OP	100,0%	100,0%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,292 <sup>a</sup>	1	,589	,719	,359	
Continuity Correction <sup>b</sup>	,130	1	,718			
Likelihood Ratio	,293	1	,588	,719	,359	
Fisher's Exact Test				,719	,359	
Linear-by-Linear Association	,291 <sup>c</sup>	1	,590	,719	,359	,124
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 18,50.

c. The standardized statistic is -,540.

## Psychische Erkrankungen vor der Schwangerschaft

Count

		Psychische Erkrankungen		Total
		Nein	Ja	
code	Mimü	108	2	110
	Vomü	104	6	110
Total		212	8	220

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,075 <sup>a</sup>	1	,150	,280	,140	
Continuity Correction <sup>b</sup>	1,167	1	,280			
Likelihood Ratio	2,168	1	,141	,280	,140	
Fisher's Exact Test				,280	,140	
Linear-by-Linear Association	2,066 <sup>c</sup>	1	,151	,280	,140	,107
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 4,00.

c. The standardized statistic is 1,437.

## Sucht

### Crosstab

Count

		Sucht		Total
		Nein	Ja	
code	Mimü	109	1	110
	vomü	108	2	110
Total		217	3	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,338 <sup>b</sup>	1	,561	1,000	,500	
Continuity Correction <sup>a</sup>	,000	1	1,000			
Likelihood Ratio	,344	1	,557	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	,336 <sup>c</sup>	1	,562	1,000	,500	,377
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,50.

c. The standardized statistic is ,580.

## Depressionen

Count

		Depressionen		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	108	2	110
Total		218	2	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,018 <sup>b</sup>	1	,155	,498	,249	
Continuity Correction <sup>a</sup>	,505	1	,477			
Likelihood Ratio	2,791	1	,095	,498	,249	
Fisher's Exact Test				,498	,249	
Linear-by-Linear Association	2,009 <sup>c</sup>	1	,156	,498	,249	,249
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is 1,417.



## Anorexie

Count

		Anorexie		Total
		Nein	Ja	
code	Mimü	109	1	110
	Vomü	110	0	110
Total		219	1	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>p</sup>	1	,316	1,000	,500	
Continuity Correction <sup>q</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is -1,000.

## Drogen

Count

		Drogen		Total
		Nein	Ja	
code	Mimü	107	3	110
	Vomü	109	1	110
Total		216	4	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,019 <sup>p</sup>	1	,313	,622	,311	
Continuity Correction <sup>q</sup>	,255	1	,614			
Likelihood Ratio	1,065	1	,302	,622	,311	
Fisher's Exact Test				,372	,311	
Linear-by-Linear Association	1,014 <sup>c</sup>	1	,314	,622	,311	,250
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,00.

c. The standardized statistic is -1,007.

## Body-Mass-Index

	code	N	Mean	Std. Deviation	Std. Error Mean
BMI	Mimü	110	21,5882	4,05795	,38691
	Vomü	108	22,8111	4,33890	,41751

### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
BMI	Equal variances assumed	,070	,791	-2,150	216	,033	-1,22293	,56887	-2,34418	-,10168
	Equal variances not assumed			-2,148	214,442	,033	-1,22293	,56922	-2,34492	-,10094

## Untergewicht

### Count

		code		Total
		Mimü	Vomü	
Untergewicht	Nein	87	98	185
	Ja	23	10	33
Total		110	108	218

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	5,757 <sup>a</sup>	1	,016	,023	,013	
Continuity Correction <sup>b</sup>	4,886	1	,027			
Likelihood Ratio	5,899	1	,015	,023	,013	
Fisher's Exact Test				,023	,013	
Linear-by-Linear Association	5,731 <sup>c</sup>	1	,017	,023	,013	,009
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 16,35.

c. The standardized statistic is -2,394.

## Anorexie nach BMI

Count		code		Total
		Mimü	Vomü	
Anorexie nach BMI	Keine Anorexie	96	103	199
	Anorextischer BMI	14	5	19
Total		110	108	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,491 <sup>a</sup>	1	,034	,053	,029	
Continuity Correction <sup>b</sup>	3,531	1	,060			
Likelihood Ratio	4,667	1	,031	,053	,029	
Fisher's Exact Test				,053	,029	
Linear-by-Linear Association	4,471 <sup>c</sup>	1	,034	,053	,029	,021
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 9,41.

c. The standardized statistic is -2,114.

## Normalgewicht

Count		code		Total
		Mimü	Vomü	
Normalgewicht	Nein	44	33	77
	Ja	66	75	141
Total		110	108	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,128 <sup>a</sup>	1	,145	,158	,094	
Continuity Correction <sup>b</sup>	1,734	1	,188			
Likelihood Ratio	2,133	1	,144	,158	,094	
Fisher's Exact Test				,158	,094	
Linear-by-Linear Association	2,118 <sup>c</sup>	1	,146	,158	,094	,039
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 38,15.

c. The standardized statistic is 1,455.

## Übergewicht

Count

		code		Total
		Mimü	Vomü	
Übergewicht	Nein	89	85	174
	Ja	21	23	44
Total		110	108	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,165 <sup>a</sup>	1	,685	,737	,406	
Continuity Correction <sup>b</sup>	,056	1	,813			
Likelihood Ratio	,165	1	,685	,737	,406	
Fisher's Exact Test				,737	,406	
Linear-by-Linear Association	,164 <sup>c</sup>	1	,686	,737	,406	,123
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 21,80.

c. The standardized statistic is ,405.

### Fehlgeburten bei früheren Schwangerschaften

Count

		Risiko Fehlgeburten		Total
		kein Risiko	Risiko	
code	Mimü	107	2	109
	Vomü	98	12	110
Total		205	14	219

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	7,534 <sup>u</sup>	1	,006	,010	,005	
Continuity Correction <sup>ß</sup>	6,093	1	,014			
Likelihood Ratio	8,316	1	,004	,010	,005	
Fisher's Exact Test				,010	,005	
Linear-by-Linear Association	7,499 <sup>c</sup>	1	,006	,010	,005	,005
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,97.

c. The standardized statistic is 2,738.

### Schwangerschaftsabbrüche in der Anamnese

Count

		Risiko Abruption		Total
		kein Risiko	Risiko	
code	Mimü	102	8	110
	Vomü	93	17	110
Total		195	25	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,655 <sup>a</sup>	1	,056	,088	,044	
Continuity Correction <sup>b</sup>	2,888	1	,089			
Likelihood Ratio	3,729	1	,053	,088	,044	
Fisher's Exact Test				,088	,044	
Linear-by-Linear Association	3,639 <sup>c</sup>	1	,056	,088	,044	,028
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 12,50.

c. The standardized statistic is 1,908.

## A3.2 Schwangerschaftsverlauf

## Ultraschalluntersuchungen und Frühdiagnostik

## Sonographie in der Schwangerschaft

	code	N	Mean	Std. Deviation	Std. Error Mean
Sonographie	Mimü	10	2,60	2,675	,846
	Vomü	28	3,36	3,009	,569

## Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Sonographie	Equal variances assumed	,009	,924	-,702	36	,487	-,757	1,079	-2,945	1,431
	Equal variances not assumed			-,743	17,762	,467	-,757	1,019	-2,901	1,386

## Count

		Sono erhalten			Total
		Ja	Nein	unbekannt	
code	Mimü	9	1	100	110
	Vomü	28	0	82	110
Total		37	1	182	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	12,537 <sup>a</sup>	2	,002	,001		
Likelihood Ratio	13,408	2	,001	,001		
Fisher's Exact Test	12,661			,001		
Linear-by-Linear Association	10,277 <sup>b</sup>	1	,001	,001	,000	,000
N of Valid Cases	220					

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,50.

b. The standardized statistic is -3,206.

## Amniozentese

Count

		Amniozentese	
		Ja	Total
code	Mimü	5	5
	Vomü	5	5
Total		10	10

## Feindiagnostik

Count

		Feindiagnostik		Total
		Nein	Ja	
code	Mimü	4	7	11
	Vomü	0	20	20
Total		4	27	31

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	8,350 <sup>a</sup>	1	,004	,010	,010	
Continuity Correction <sup>b</sup>	5,428	1	,020			
Likelihood Ratio	9,421	1	,002	,010	,010	
Fisher's Exact Test				,010	,010	
Linear-by-Linear Association	8,081 <sup>c</sup>	1	,004	,010	,010	,010
N of Valid Cases	31					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,42.

c. The standardized statistic is 2,843.



## Medikamentöse Prophylaxe

Count

		Prophylaxe		Total
		Keine Prophylaxe	Prophylaxe	
code	Mimü	17	56	73
	Vomü	16	76	92
Total		33	132	165

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,884 <sup>a</sup>	1	,347	,434	,228	
Continuity Correction <sup>b</sup>	,554	1	,457			
Likelihood Ratio	,880	1	,348	,434	,228	
Fisher's Exact Test				,434	,228	
Linear-by-Linear Association	,879 <sup>c</sup>	1	,348	,434	,228	,100
N of Valid Cases	165					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,60.

c. The standardized statistic is ,938.

## Sexuell übertragene Infektionen

STD gesamt

code		N	Mean	Std. Deviation	Std. Error Mean
STD	Mimü	109	,24	,489	,047
	Vomü	109	,14	,440	,042

### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
								Lower	Upper	
STD	Equal variances assumed	8,220	,005	1,602	216	,111	,101	,063	-,023	,225
	Equal variances not assumed			1,602	213,687	,111	,101	,063	-,023	,225

## Gesamtanzahl infizierter Frauen

Count

		STD		Total
		Nein	Ja	
code	Mimü	86	23	109
	Vomü	97	12	109
Total		183	35	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,118 <sup>u</sup>	1	,042	,064	,032	
Continuity Correction <sup>ß</sup>	3,404	1	,065			
Likelihood Ratio	4,178	1	,041	,064	,032	
Fisher's Exact Test				,064	,032	
Linear-by-Linear Association	4,099 <sup>c</sup>	1	,043	,064	,032	,019
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 17,50.

c. The standardized statistic is -2,025.

## Clamydien

Count

		Clamydien		Total
		Nein	Ja	
code	Mimü	98	11	109
	Vomü	104	5	109
Total		202	16	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,428 <sup>a</sup>	1	,119	,193	,096	
Continuity Correction <sup>b</sup>	1,686	1	,194			
Likelihood Ratio	2,484	1	,115	,193	,096	
Fisher's Exact Test				,193	,096	
Linear-by-Linear Association	2,417 <sup>c</sup>	1	,120	,193	,096	,063
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 8,00.

c. The standardized statistic is -1,555.

## Humanes Papilom Virus

Count

		HPV		Total
		Nein	Ja	
code	Mimü	108	1	109
	Vomü	109	0	109
Total		217	1	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is -1,000.

## Hepatitis-B Virus

Count

		HBV		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	108	1	109
Total		217	1	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is -1,000.

## Hepatitis-C Virus

Count

		HCV		Total
		Nein	Ja	
code	Mimü	108	1	109
	Vomü	109	0	109
Total		217	1	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is -1,000.

## Pilze

Count

		Pilze		Total
		Nein	Ja	
code	Mimü	99	10	109
	Vomü	101	8	109
Total		200	18	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,242 <sup>a</sup>	1	,623	,806	,403	
Continuity Correction <sup>b</sup>	,061	1	,806			
Likelihood Ratio	,243	1	,622	,806	,403	
Fisher's Exact Test				,806	,403	
Linear-by-Linear Association	,241 <sup>c</sup>	1	,623	,806	,403	,173
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 9,00.

c. The standardized statistic is -,491.

## B-Streptokokken

Count

		B Streptokokken		Total
		Nein	Ja	
code	Mimü	106	3	109
	Vomü	108	1	109
Total		214	4	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,019 <sup>a</sup>	1	,313	,622	,311	
Continuity Correction <sup>b</sup>	,255	1	,614			
Likelihood Ratio	1,065	1	,302	,622	,311	
Fisher's Exact Test				,622	,311	
Linear-by-Linear Association	1,014 <sup>c</sup>	1	,314	,622	,311	,250
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,00.

c. The standardized statistic is -1,007.

## Weitere Infektionen

Infektionen gesamt

code		N	Mean	Std. Deviation	Std. Error Mean
Infekte	Mimü	109	,09	,320	,031
	Vomü	109	,17	,381	,037

## Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Infekte	Equal variances assumed	11,478	,001	-1,731	216	,085	-,083	,048	-,177	,011
	Equal variances not assumed			-1,731	209,791	,085	-,083	,048	-,177	,011

## Gesamtanzahl betroffener Frauen

## Count

		Infekte		Total
		Nein	Ja	
code	Mimü	100	9	109
	Vomü	90	19	109
Total		190	28	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,098 <sup>a</sup>	1	,043	,067	,034	
Continuity Correction <sup>b</sup>	3,319	1	,068			
Likelihood Ratio	4,178	1	,041	,067	,034	
Fisher's Exact Test				,067	,034	
Linear-by-Linear Association	4,079 <sup>c</sup>	1	,043	,067	,034	,021
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,00.

c. The standardized statistic is 2,020.

## Atemwegsinfekte

## Count

		Atemwegsinfekte		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	105	5	110
Total		214	5	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	5,070 <sup>a</sup>	1	,024	,060	,031	
Continuity Correction <sup>b</sup>	3,238	1	,072			
Likelihood Ratio	7,002	1	,008	,060	,031	
Fisher's Exact Test				,060	,031	
Linear-by-Linear Association	5,047 <sup>c</sup>	1	,025	,060	,031	,031
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,49.

c. The standardized statistic is 2,247.

## Harnwegsinfekte

## Count

		HWI		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	107	2	109
Total		216	2	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,019 <sup>a</sup>	1	,155	,498	,249	
Continuity Correction <sup>b</sup>	,505	1	,477			
Likelihood Ratio	2,791	1	,095	,498	,249	
Fisher's Exact Test				,498	,249	
Linear-by-Linear Association	2,009 <sup>c</sup>	1	,156	,498	,249	,249
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is 1,417.

## Pyelonephritiden

## Count

		Pyelonephritiden		Total
		Nein	Ja	
code	Mimü	106	3	109
	Vomü	110	0	110
Total		216	3	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,070 <sup>a</sup>	1	,080	,122	,122	
Continuity Correction <sup>b</sup>	1,370	1	,242			
Likelihood Ratio	4,228	1	,040	,122	,122	
Fisher's Exact Test				,122	,122	
Linear-by-Linear Association	3,056 <sup>c</sup>	1	,080	,122	,122	,122
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,49.

c. The standardized statistic is -1,748.

## Magen-Darminfektionen

Count

		Magendarminfektionen		Total
		Nein	Ja	
code	Mimü	107	2	109
	Vomü	110	0	110
Total		217	2	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,037 <sup>a</sup>	1	,154	,247	,247	
Continuity Correction <sup>b</sup>	,514	1	,473			
Likelihood Ratio	2,809	1	,094	,247	,247	
Fisher's Exact Test				,247	,247	
Linear-by-Linear Association	2,028 <sup>c</sup>	1	,154	,247	,247	,247
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is -1,424.

## Windpocken/ Gürtelrose

Count

		Windpocken/ Gürtelrose	Total
		Nein	
code	Mimü	109	109
	Vomü	110	110
Total		219	219



## Röteln

Count

		Röteln		Total
		Nein	Ja	
code	Mimü	108	1	109
	Vomü	109	0	109
Total		217	1	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is -1,000.

## Erythema infectiosum

Count

		Erythema infectiosum	
		Nein	Total
code	Mimü	109	109
	Vomü	110	110
Total		219	219

## Toxoplasmose

Count

		Toxoplasmose		Total
		Nein	Ja	
code	Mimü	106	3	109
	Vomü	109	0	109
Total		215	3	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,042 <sup>a</sup>	1	,081	,247	,123	
Continuity Correction <sup>b</sup>	1,352	1	,245			
Likelihood Ratio	4,201	1	,040	,247	,123	
Fisher's Exact Test				,247	,123	
Linear-by-Linear Association	3,028 <sup>c</sup>	1	,082	,247	,123	,123
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,50.

c. The standardized statistic is -1,740.

## andere Infektionen

Count

		andere Infektionen		Total
		Nein	Ja	
code	Mimü	105	3	108
	Vomü	97	13	110
Total		202	16	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	6,549 <sup>a</sup>	1	,010	,017	,009	
Continuity Correction <sup>b</sup>	5,287	1	,021			
Likelihood Ratio	7,037	1	,008	,017	,009	
Fisher's Exact Test				,017	,009	
Linear-by-Linear Association	6,519 <sup>c</sup>	1	,011	,017	,009	,008
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,93.

c. The standardized statistic is 2,553.

## Schwangerschaftserkrankungen

## Schwangerschaftserkrankungen gesamt

	code	N	Mean	Std. Deviation	Std. Error Mean
SSErkrankungen	Mimü	110	,03	,164	,016
	Vomü	110	,12	,351	,034

## Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
SS Erkrankungen	Equal variances assumed	26,576	,000	-2,459	218	,015	-,091	,037	-,164	-,018
	Equal variances not assumed			-2,459	154,130	,015	-,091	,037	-,164	-,018

## Gesamtanzahl betroffener Frauen

## Count

		SS Erkrankungen		Total
		Nein	Ja	
code	Mimü	107	3	110
	Vomü	98	12	110
Total		205	15	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	5,795 <sup>a</sup>	1	,016	,029	,015	
Continuity Correction <sup>b</sup>	4,579	1	,032			
Likelihood Ratio	6,178	1	,013	,029	,015	
Fisher's Exact Test				,029	,015	
Linear-by-Linear Association	5,769 <sup>c</sup>	1	,016	,029	,015	,012
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,50.

c. The standardized statistic is 2,402.

## Emesis

## Count

		Emesis		Total
		Nein	Ja	
code	Mimü	109	1	110
	Vomü	105	5	110
Total		214	6	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,741 <sup>d</sup>	1	,098	,212	,106	
Continuity Correction <sup>e</sup>	1,542	1	,214			
Likelihood Ratio	2,986	1	,084	,212	,106	
Fisher's Exact Test				,212	,106	
Linear-by-Linear Association	2,729 <sup>c</sup>	1	,099	,212	,106	,092
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 3,00.

c. The standardized statistic is 1,652.

## isolierte Hypertonie

## Count

		isolierte Hypertonie		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	105	5	110
Total		215	5	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	5,116 <sup>d</sup>	1	,024	,060	,030	
Continuity Correction <sup>e</sup>	3,274	1	,070			
Likelihood Ratio	7,048	1	,008	,060	,030	
Fisher's Exact Test				,060	,030	
Linear-by-Linear Association	5,093 <sup>c</sup>	1	,024	,060	,030	,030
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,50.

c. The standardized statistic is 2,257.

## Präeklampsie

Count

		Präeklampsie		Total
		Nein	Ja	
code	Mimü	108	2	110
	Vomü	109	1	110
Total		217	3	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,338 <sup>a</sup>	1	,561	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,344	1	,557	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	,336 <sup>c</sup>	1	,562	1,000	,500	,377
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,50.

c. The standardized statistic is -,580.

## Eklampsie

Count

		Eklampsie	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## HELLP-Syndrom

Count

		HELLP		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	108	2	110
Total		218	2	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,018 <sup>a</sup>	1	,155	,498	,249	
Continuity Correction <sup>b</sup>	,505	1	,477			
Likelihood Ratio	2,791	1	,095	,498	,249	
Fisher's Exact Test				,498	,249	
Linear-by-Linear Association	2,009 <sup>c</sup>	1	,156	,498	,249	,249
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is 1,417.

## Neu aufgetretene Erkrankungen während der Schwangerschaft

### Erkrankungen in der Schwangerschaft gesamt

	code	N	Mean	Std. Deviation	Std. Error Mean
Erkrankungen in der Schwangerschaft	Mimü	110	,08	,275	,026
	Vomü	110	,18	,410	,039

#### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Erkrankungen in der Schwangerschaft	Equal variances assumed	19,159	,000	-2,122	218	,035	-,100	,047	-,193	-,007
	Equal variances not assumed			-2,122	190,580	,035	-,100	,047	-,193	-,007

### Gesamtanzahl erkrankter Frauen

#### Count

		Erkrankungen während der Schwangerschaft		Total
		Nein	Ja	
code	Mimü	103	7	110
	Vomü	92	17	109
Total		195	24	219

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,783 <sup>d</sup>	1	,029	,032	,024	
Continuity Correction <sup>e</sup>	3,883	1	,049			
Likelihood Ratio	4,913	1	,027	,032	,024	
Fisher's Exact Test				,032	,024	
Linear-by-Linear Association	4,761 <sup>c</sup>	1	,029	,032	,024	,016
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 11,95.

c. The standardized statistic is 2,182.

## Herz-Kreislauf-Erkrankungen in der Schwangerschaft

Count

		Herz Kreislauf Erkrankungen		Total
		Nein	Ja	
code	Mimü	107	3	110
	Vomü	105	5	110
Total		212	8	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,519 <sup>a</sup>	1	,471	,721	,361	
Continuity Correction <sup>b</sup>	,130	1	,719			
Likelihood Ratio	,524	1	,469	,721	,361	
Fisher's Exact Test				,721	,361	
Linear-by-Linear Association	,517 <sup>c</sup>	1	,472	,721	,361	,221
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 4,00.

c. The standardized statistic is ,719.

## chronische Gelenkerkrankungen in der Schwangerschaft

Count

		chronische Gelenkerkrankungen		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	110	0	110
Total		220	0	220

## chronische Lungenerkrankungen in der Schwangerschaft

Count

		chronische Lungenerkrankungen		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	110	0	110
Total		220	0	220



---

 chronische Magen-Darm-Erkrankungen in der Schwangerschaft

Count

		Chronische Magen-Damerkrankungen	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## Geschlechtskrankheiten in der Schwangerschaft

Count

		Geschlechtskrankheiten	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## Stoffwechselerkrankungen in der Schwangerschaft

Count

		Stoffwechselerkrankungen		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	109	1	110
Total		219	1	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is 1,000.

## Diabetes in der Schwangerschaft

Count

		Diabetes		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	109	1	110
Total		219	1	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is 1,000.

## chronische Schmerzen in der Schwangerschaft

Count

		chronische Schmerzen	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## Migräne in der Schwangerschaft

Count

		Migräne	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## Gynäkologische Erkrankungen in der Schwangerschaft

Count

		Gynäkologische Erkrankungen		Total
		Nein	Ja	
code	Mimü	109	1	110
	Vomü	109	1	110
Total		218	2	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,000 <sup>a</sup>	1	1,000	1,000	,751	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,000	1	1,000	1,000	,751	
Fisher's Exact Test				1,000	,751	
Linear-by-Linear Association	,000 <sup>c</sup>	1	1,000	1,000	,751	,502
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is ,000.

## Leber- / Gallen- Erkrankungen in der Schwangerschaft

Count

		Leber oder Gallenerkrankungen		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	109	1	110
Total		219	1	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is 1,000.

## Allergien in der Schwangerschaft

Count

		Allergien	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## Tumor in der Schwangerschaft

Count

		Tumor	
		Nein	Total
code	Mimü	110	110
	Vomü	110	110
Total		220	220

## andere Erkrankungen in der Schwangerschaft

Count

		andere Erkrankungen		Total
		Nein	Ja	
code	Mimü	105	5	110
	Vomü	99	11	110
Total		204	16	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,426 <sup>a</sup>	1	,119	,193	,097	
Continuity Correction <sup>b</sup>	1,685	1	,194			
Likelihood Ratio	2,482	1	,115	,193	,097	
Fisher's Exact Test				,193	,097	
Linear-by-Linear Association	2,415 <sup>c</sup>	1	,120	,193	,097	,063
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 8,00.

c. The standardized statistic is 1,554.

## Weitere Probleme während des Schwangerschaftsverlauf

### Zervixinsuffizienz

Count

		Zervixinsuffizienz		Total
		Nein	Ja	
code	Mimü	104	6	110
	Vomü	104	6	110
Total		208	12	220

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,000 <sup>a</sup>	1	1,000	1,000	,616	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,000	1	1,000	1,000	,616	
Fisher's Exact Test				1,000	,616	
Linear-by-Linear Association	,000 <sup>c</sup>	1	1,000	1,000	,616	,232
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,00.

c. The standardized statistic is ,000.

## Plazentainsuffizienz in der Schwangerschaft

Count

		Plazentainsuffizienz		Total
		Nein	Ja	
code	Mimü	106	4	110
	Vomü	107	3	110
Total		213	7	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,148 <sup>a</sup>	1	,701	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,148	1	,700	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	,147 <sup>c</sup>	1	,702	1,000	,500	,277
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 3,50.

c. The standardized statistic is -,383.

## Amnioninfektionssyndrom

Count

		Amnioninfektionssyndrom		Total
		Nein	Ja	
code	Mimü	108	2	110
	Vomü	107	3	110
Total		215	5	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,205 <sup>a</sup>	1	,651	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,206	1	,650	1,000	,500	
Fisher's Exact Test				,685	,500	
Linear-by-Linear Association	,204 <sup>c</sup>	1	,652	1,000	,500	,315
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,50.

c. The standardized statistic is ,451.

## Blutung

Count

		Blutung		Total
		Nein	Ja	
code	Mimü	108	2	110
	Vomü	105	5	110
Total		213	7	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,328 <sup>a</sup>	1	,249	,446	,223	
Continuity Correction <sup>b</sup>	,590	1	,442			
Likelihood Ratio	1,371	1	,242	,446	,223	
Fisher's Exact Test				,446	,223	
Linear-by-Linear Association	1,322 <sup>c</sup>	1	,250	,446	,223	,163
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 3,50.

c. The standardized statistic is 1,150.

## Chirurgische Eingriffe während der Schwangerschaft

Count

		Chirurgische Eingriffe		Total
		Nein	Ja	
code	Mimü	108	1	109
	Vomü	107	3	110
Total		215	4	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,000 <sup>a</sup>	1	,317	,622	,314	
Continuity Correction <sup>b</sup>	,245	1	,620			
Likelihood Ratio	1,047	1	,306	,622	,314	
Fisher's Exact Test				,622	,314	
Linear-by-Linear Association	,996 <sup>c</sup>	1	,318	,622	,314	,252
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,99.

c. The standardized statistic is ,998.

## Unfälle während der Schwangerschaft

Count

		Unfälle		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	107	3	110
Total		216	3	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,014 <sup>p</sup>	1	,083	,247	,125	
Continuity Correction <sup>q</sup>	1,333	1	,248			
Likelihood Ratio	4,173	1	,041	,247	,125	
Fisher's Exact Test				,247	,125	
Linear-by-Linear Association	3,000 <sup>c</sup>	1	,083	,247	,125	,125
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,49.

c. The standardized statistic is 1,732.

## Lungenreife der Mutter erfolgt

Count

		Lungenreife Mutter		Total
		Nein	Ja	
code	Mimü	96	13	109
	Vomü	93	15	108
Total		189	28	217

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,186 <sup>p</sup>	1	,666	,691	,410	
Continuity Correction <sup>q</sup>	,052	1	,819			
Likelihood Ratio	,186	1	,666	,691	,410	
Fisher's Exact Test				,691	,410	
Linear-by-Linear Association	,185 <sup>c</sup>	1	,667	,691	,410	,146
N of Valid Cases	217					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 13,94.

c. The standardized statistic is ,430.



### Stationärer Aufenthalt

	code	N	Mean	Std. Deviation	Std. Error Mean
Stationärer Aufenthalt	Mimü	41	1,90	1,947	,304
	Vomü	39	15,00	18,121	2,902

#### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Stationärer Aufenthalt	Equal variances assumed	29,347	,000	-4,602	78	,000	-13,098	2,846	-18,764	-7,431
	Equal variances not assumed			-4,489	38,835	,000	-13,098	2,918	-19,000	-7,195

#### Count

		Krankenhausaufenthalt während der Schwangerschaft		Total
		Nein	Ja	
code	Mimü	66	44	110
	Vomü	70	39	109
Total		136	83	219

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,414 <sup>a</sup>	1	,520	,578	,307	
Continuity Correction <sup>b</sup>	,254	1	,614			
Likelihood Ratio	,414	1	,520	,578	,307	
Fisher's Exact Test				,578	,307	
Linear-by-Linear Association	,412 <sup>c</sup>	1	,521	,578	,307	,090
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 41,31.

c. The standardized statistic is -,642.

## Gewichtszunahme während der Schwangerschaft

	code	N	Mean	Std. Deviation	Std. Error Mean
Gewicht zur Geburt	Mimü	103	14,047	6,4781	,6383
	Vomü	99	14,378	7,1464	,7182

### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Gewicht zur Geburt	Equal variances assumed	,412	,522	-,345	200	,730	-,3312	,9590	-2,2223	1,5599
	Equal variances not assumed			-,345	196,290	,731	-,3312	,9609	-2,2262	1,5638

## A3.3 Geburt

### Schwangerschaftsdauer

Schwangerschaftsdauer in Tagen

entbWoch

Mimü	N	Valid	102
		Missing	9
	Mean		272,11
	Median		277,50
	Std. Deviation		18,839
	Minimum		178
	Maximum		294
Vomü	N	Valid	110
		Missing	0
	Mean		272,70
	Median		279,00
	Std. Deviation		19,668
	Minimum		185
	Maximum		293

## Frühgeburt

Count

		Frühgeburt		Total
		Ja	Nein	
code	Mimü	16	86	102
	Vomü	14	96	110
Total		30	182	212

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,381 <sup>u</sup>	1	,537	,560	,337	
Continuity Correction <sup>b</sup>	,177	1	,674			
Likelihood Ratio	,381	1	,537	,560	,337	
Fisher's Exact Test				,560	,337	
Linear-by-Linear Association	,380 <sup>c</sup>	1	,538	,560	,337	,129
N of Valid Cases	212					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,43.

c. The standardized statistic is ,616.

## Art der Aufnahme zur Geburt

			Aufnahme zur Geburt		Total
			Normale Aufnahme	Notfall	
code	Mimü	Count	84	26	110
		Expected Count	91,0	19,0	110,0
		% within code	76,4%	23,6%	100,0%
		% within Aufn_Typ	46,2%	68,4%	50,0%
	Vomü	Count	98	12	110
		Expected Count	91,0	19,0	110,0
		% within code	89,1%	10,9%	100,0%
		% within Aufn_Typ	53,8%	31,6%	50,0%
Total	Count	182	38	220	
	Expected Count	182,0	38,0	220,0	
	% within code	82,7%	17,3%	100,0%	
	% within Aufn_Typ	100,0%	100,0%	100,0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	6,235 <sup>a</sup>	1	,013	,020	,010	
Continuity Correction <sup>b</sup>	5,376	1	,020			
Likelihood Ratio	6,359	1	,012	,020	,010	
Fisher's Exact Test				,020	,010	
Linear-by-Linear Association	6,206 <sup>c</sup>	1	,013	,020	,010	,006
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 19,00.

c. The standardized statistic is -2,491.

## Entbindungsverlauf

## Geburtsbeginn

		Spontaner Geburtsbeginn		Total
		Nein	Ja	
code	Mimü	35	62	97
	Vomü	19	85	104
Total		54	147	201

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	8,105 <sup>a</sup>	1	,004	,007	,003	
Continuity Correction <sup>b</sup>	7,224	1	,007			
Likelihood Ratio	8,182	1	,004	,007	,003	
Fisher's Exact Test				,007	,003	
Linear-by-Linear Association	8,065 <sup>c</sup>	1	,005	,007	,003	,002
N of Valid Cases	201					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 26,06.

c. The standardized statistic is 2,840.

## Count

		Priming/ Wehenfördernde Mittel		Total
		Nein	Ja	
code	Mimü	62	35	97
	Vomü	85	19	104
Total		147	54	201

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	8,105 <sup>a</sup>	1	,004	,007	,003	
Continuity Correction <sup>b</sup>	7,224	1	,007			
Likelihood Ratio	8,182	1	,004	,007	,003	
Fisher's Exact Test				,007	,003	
Linear-by-Linear Association	8,065 <sup>c</sup>	1	,005	,007	,003	,002
N of Valid Cases	201					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 26,06.

c. The standardized statistic is -2,840.

## Spontanentbindung

Count

		Spontanentbindeung		Total
		Nein	Ja	
code	Mimü	21	88	109
	Vomü	28	82	110
Total		49	170	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,207 <sup>a</sup>	1	,272	,331	,175	
Continuity Correction <sup>b</sup>	,877	1	,349			
Likelihood Ratio	1,211	1	,271	,331	,175	
Fisher's Exact Test				,331	,175	
Linear-by-Linear Association	1,202 <sup>c</sup>	1	,273	,331	,175	,071
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 24,39.

c. The standardized statistic is -1,096.

## Sectio caesera

Count

		Sectio Gesamt		Total
		Nein	Ja	
code	Mimü	93	17	110
	Vomü	88	22	110
Total		181	39	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,779 <sup>a</sup>	1	,377		
Continuity Correction <sup>b</sup>	,499	1	,480		
Likelihood Ratio	,781	1	,377		
Fisher's Exact Test				,480	,240
Linear-by-Linear Association	,776	1	,378		
N of Valid Cases	220				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 19,50.

## primäre Sectio

Count

		primäre Sektio		Total
		Nein	Ja	
code	Mimü	99	11	110
	Vomü	95	15	110
Total		194	26	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,698 <sup>a</sup>	1	,404	,532	,266	
Continuity Correction <sup>b</sup>	,393	1	,531			
Likelihood Ratio	,700	1	,403	,532	,266	
Fisher's Exact Test				,532	,266	
Linear-by-Linear Association	,695 <sup>c</sup>	1	,405	,532	,266	,118
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 13,00.

c. The standardized statistic is ,833.

## sekundäre Sectio

Count

		sekundäre Sectio		Total
		Nein	Ja	
code	Mimü	104	6	110
	Vomü	103	7	110
Total		207	13	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,082 <sup>a</sup>	1	,775	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,082	1	,775	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	,081 <sup>c</sup>	1	,775	1,000	,500	,215
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,50.

c. The standardized statistic is ,285.

## Vakuumextraktion

Count

		Vakuumextraktion		Total
		Nein	Ja	
code	Mimü	108	1	109
	Vomü	109	1	110
Total		217	2	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,000 <sup>a</sup>	1	,995	1,000	,749	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,000	1	,995	1,000	,749	
Fisher's Exact Test				1,000	,749	
Linear-by-Linear Association	,000 <sup>c</sup>	1	,995	1,000	,749	,502
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is -,006.



## Forceps

Count

		Forceps		Total
		Nein	Ja	
code	Mimü	106	3	109
	Vomü	103	7	110
Total		209	10	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,639 <sup>a</sup>	1	,201	,332	,170	
Continuity Correction <sup>b</sup>	,915	1	,339			
Likelihood Ratio	1,684	1	,194	,332	,170	
Fisher's Exact Test				,332	,170	
Linear-by-Linear Association	1,631 <sup>c</sup>	1	,202	,332	,170	,118
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,98.

c. The standardized statistic is 1,277.

## Schädellage

Count

		Schädellage		Total
		Nein	Ja	
code	Mimü	5	91	96
	Vomü	11	95	106
Total		16	186	202

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,845 <sup>a</sup>	1	,174	,201	,136	
Continuity Correction <sup>b</sup>	1,205	1	,272			
Likelihood Ratio	1,897	1	,168	,201	,136	
Fisher's Exact Test				,201	,136	
Linear-by-Linear Association	1,836 <sup>c</sup>	1	,175	,201	,136	,085
N of Valid Cases	202					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,60.

c. The standardized statistic is -1,355.

## Beckenlage

Count		Beckenlage		Total
		Nein	Ja	
code	Mimü	93	3	96
	Vomü	99	7	106
Total		192	10	202

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,296 <sup>a</sup>	1	,255	,338	,209	
Continuity Correction <sup>b</sup>	,662	1	,416			
Likelihood Ratio	1,338	1	,247	,338	,209	
Fisher's Exact Test				,338	,209	
Linear-by-Linear Association	1,289 <sup>c</sup>	1	,256	,338	,209	,140
N of Valid Cases	202					

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,75.

c. The standardized statistic is 1,135.

## Querlage

Count		Querlage		Total
		Nein	Ja	
code	Mimü	94	2	96
	Vomü	102	4	106
Total		196	6	202

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,499 <sup>a</sup>	1	,480	,685	,389	
Continuity Correction <sup>b</sup>	,085	1	,771			
Likelihood Ratio	,511	1	,475	,685	,389	
Fisher's Exact Test				,685	,389	
Linear-by-Linear Association	,497 <sup>c</sup>	1	,481	,685	,389	,259
N of Valid Cases	202					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,85.

c. The standardized statistic is ,705.

## Blasensprung

## Wann war der Blasensprung

code			Frequency	Percent	Valid Percent	Cumulative Percent
Mimü	Valid	unter 4h vor Geburt	49	44,1	49,5	49,5
		zw 4-12 h vor Geburt	37	33,3	37,4	86,9
		über 12h vor Geburt	11	9,9	11,1	98,0
		unter der Geburt	2	1,8	2,0	100,0
		Total	99	89,2	100,0	
	Missing	88	3	2,7		
		unbekannt	8	7,2		
		System	1	,9		
		Total	12	10,8		
		Total	111	100,0		
Vomü	Valid	unter 4h vor Geburt	36	32,7	37,9	37,9
		zw 4-12 h vor Geburt	40	36,4	42,1	80,0
		über 12h vor Geburt	18	16,4	18,9	98,9
		unter der Geburt	1	,9	1,1	100,0
		Total	95	86,4	100,0	
	Missing	unbekannt	15	13,6		
	Total	110	100,0			

## Count

		Blasensprung		Total
		spontan	künstlich	
code	Mimü	41	59	100
	Vomü	54	48	102
Total		95	107	202

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,890 <sup>a</sup>	1	,089	,093	,059	
Continuity Correction <sup>b</sup>	2,431	1	,119			
Likelihood Ratio	2,898	1	,089	,093	,059	
Fisher's Exact Test				,093	,059	
Linear-by-Linear Association	2,876 <sup>c</sup>	1	,090	,093	,059	,027
N of Valid Cases	202					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 47,03.

c. The standardized statistic is -1,696.

Count

		spontaner Blasensprung		Total
		Nein	Ja	
code	Mimü	59	41	100
	Vomü	48	54	102
Total		107	95	202

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,890 <sup>a</sup>	1	,089	,093	,059	
Continuity Correction <sup>b</sup>	2,431	1	,119			
Likelihood Ratio	2,898	1	,089	,093	,059	
Fisher's Exact Test				,093	,059	
Linear-by-Linear Association	2,876 <sup>c</sup>	1	,090	,093	,059	,027
N of Valid Cases	202					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 47,03.

c. The standardized statistic is 1,696.

Count

		künstlicher Blasensprung		Total
		Nein	Ja	
code	Mimü	41	59	100
	Vomü	54	48	102
Total		95	107	202

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,890 <sup>a</sup>	1	,089	,093	,059	
Continuity Correction <sup>b</sup>	2,431	1	,119			
Likelihood Ratio	2,898	1	,089	,093	,059	
Fisher's Exact Test				,093	,059	
Linear-by-Linear Association	2,876 <sup>c</sup>	1	,090	,093	,059	,027
N of Valid Cases	202					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 47,03.

c. The standardized statistic is -1,696.

## Pathologischer Blasensprung

Count

		pathologischer Blasensprung		Total
		Nein	Ja	
code	Mimü	88	11	99
	Vomü	77	18	95
Total		165	29	194

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,342 <sup>d</sup>	1	,126	,159	,092	
Continuity Correction <sup>e</sup>	1,766	1	,184			
Likelihood Ratio	2,358	1	,125	,159	,092	
Fisher's Exact Test				,159	,092	
Linear-by-Linear Association	2,329 <sup>c</sup>	1	,127	,159	,092	,051
N of Valid Cases	194					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,20.

c. The standardized statistic is 1,526.

## Pathologisches Fruchtwasser

Count

		pathologisches Fruchtwasser		Total
		Nein	Ja	
code	Mimü	97	13	110
	Vomü	95	15	110
Total		192	28	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,164 <sup>d</sup>	1	,686	,840	,420	
Continuity Correction <sup>e</sup>	,041	1	,840			
Likelihood Ratio	,164	1	,686	,840	,420	
Fisher's Exact Test				,840	,420	
Linear-by-Linear Association	,163 <sup>c</sup>	1	,686	,840	,420	,148
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,00.

c. The standardized statistic is ,404.

## Blutverlust unter der Geburt

Count

		Blutverlust		Total
		normaler Blutverlust	erhöhter Blutverlust	
code	Mimü	97	7	104
	Vomü	98	11	109
Total		195	18	213

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,777 <sup>a</sup>	1	,378	,463	,264	
Continuity Correction <sup>b</sup>	,403	1	,525			
Likelihood Ratio	,784	1	,376	,463	,264	
Fisher's Exact Test				,463	,264	
Linear-by-Linear Association	,773 <sup>c</sup>	1	,379	,463	,264	,134
N of Valid Cases	213					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 8,79.

c. The standardized statistic is ,879.

## Dauer der Geburt in Minuten

Dauer

Mimü	N	Valid	93
		Missing	18
	Mean		410,53
	Median		375,00
	Std. Deviation		185,825
	Minimum		60
	Maximum		999
Vomü	N	Valid	107
		Missing	3
	Mean		438,18
	Median		420,00
	Std. Deviation		322,431
	Minimum		20
	Maximum		2760

## Komplikationen unter der Geburt

### Geburtskomplikationen gesamt

	code	N	Mean	Std. Deviation	Std. Error Mean
Komplikationen unter der Geburt	Mimü	109	,24	,489	,047
	Vomü	110	,81	,748	,071

#### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Komplikationen unter der Geburt	Equal variances assumed	23,595	,000	-6,676	217	,000	-,571	,085	-,739	-,402
	Equal variances not assumed			-6,688	187,943	,000	-,571	,085	-,739	-,402

### Gesamtanzahl betroffener Frauen

#### Count

		Komplikationen unter der Geburt		Total
		Nein	Ja	
code	Mimü	86	23	109
	Vomü	42	68	110
Total		128	91	219

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	37,374 <sup>a</sup>	1	,000	,000	,000	
Continuity Correction <sup>b</sup>	35,716	1	,000			
Likelihood Ratio	38,695	1	,000	,000	,000	
Fisher's Exact Test				,000	,000	
Linear-by-Linear Association	37,203 <sup>c</sup>	1	,000	,000	,000	,000
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 45,29.

c. The standardized statistic is 6,099.

## Plazenta praevia

Count

		Plazenta praevia		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	108	2	110
Total		217	2	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,000 <sup>a</sup>	1	,157	,498	,251	
Continuity Correction <sup>b</sup>	,495	1	,482			
Likelihood Ratio	2,773	1	,096	,498	,251	
Fisher's Exact Test				,498	,251	
Linear-by-Linear Association	1,991 <sup>c</sup>	1	,158	,498	,251	,251
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is 1,411.

## tiefer Sitz der Plazenta

Count

		tiefer Sitz der Plazenta	
		Nein	Total
code	Mimü	109	109
	Vomü	110	110
Total		219	219



## vorzeitige Plazentalösung

Count

		Plazentaablösung		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	108	2	110
Total		217	2	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,000 <sup>a</sup>	1	,157	,498	,251	
Continuity Correction <sup>b</sup>	,495	1	,482			
Likelihood Ratio	2,773	1	,096	,498	,251	
Fisher's Exact Test				,498	,251	
Linear-by-Linear Association	1,991 <sup>c</sup>	1	,158	,498	,251	,251
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is 1,411.

## Randsinusblutung

Count

		Randsinusblutung		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	110	0	110
Total		219	0	219

## Uterusruptur

Count

		Uterusruptur		Total
		Nein	Ja	
code	Mimü	109	0	109
	Vomü	110	0	110
Total		219	0	219

## Gefäßruptur bei Insertio Velamentosa

Count

		Gefäßruptur	
		Nein	Total
code	Mimü	109	109
	Vomü	110	110
Total		219	219

## Varixknotenruptur der Scheide

Count

		Varixknotenruptur	
		Nein	Total
code	Mimü	109	109
	Vomü	110	110
Total		219	219

## Scheidenriss

Count

		Scheidenriss		Total
		Nein	Ja	
code	Mimü	93	16	109
	Vomü	92	18	110
Total		185	34	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,118 <sup>d</sup>	1	,731	,852	,438	
Continuity Correction <sup>e</sup>	,025	1	,875			
Likelihood Ratio	,119	1	,731	,852	,438	
Fisher's Exact Test				,852	,438	
Linear-by-Linear Association	,118 <sup>c</sup>	1	,731	,852	,438	,139
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 16,92.

c. The standardized statistic is ,343.

## Zervixriss

Count

		Zervixriss		Total
		Nein	Ja	
code	Mimü	108	1	109
	Vomü	110	0	110
Total		218	1	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,014 <sup>d</sup>	1	,314	,498	,498	
Continuity Correction <sup>e</sup>	,000	1	,996			
Likelihood Ratio	1,400	1	,237	,498	,498	
Fisher's Exact Test				,498	,498	
Linear-by-Linear Association	1,009 <sup>c</sup>	1	,315	,498	,498	,498
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is -1,005.

## Schock

Count

		Schock	
		Nein	Total
code	Mimü	109	109
	Vomü	110	110
Total		219	219

## Atonie

Count

		Atonie		Total
		Nein	Ja	
code	Mimü	107	2	109
	Vomü	108	2	110
Total		215	4	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,000 <sup>a</sup>	1	,993	1,000	,686	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,000	1	,993	1,000	,686	
Fisher's Exact Test				1,000	,686	
Linear-by-Linear Association	,000 <sup>c</sup>	1	,993	1,000	,686	,378
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,99.

c. The standardized statistic is -,009.

## andere Komplikationen

Count

		andere Komplikationen		Total
		Nein	Ja	
code	Mimü	104	5	109
	Vomü	59	51	110
Total		163	56	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	50,206 <sup>a</sup>	1	,000	,000	,000	
Continuity Correction <sup>b</sup>	48,034	1	,000			
Likelihood Ratio	56,515	1	,000	,000	,000	
Fisher's Exact Test				,000	,000	
Linear-by-Linear Association	49,976 <sup>c</sup>	1	,000	,000	,000	,000
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 27,87.

c. The standardized statistic is 7,069.

## Komplikationen im Wochenbett

### Komplikationen im Wochenbett gesamt

	code	N	Mean	Std. Deviation	Std. Error Mean
Wochenbett-komplikationen	Mimü	105	,46	,519	,051
	Vomü	110	,59	,547	,052

#### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Wochenbett-komplikationen	Equal variances assumed	,282	,596	-1,837	213	,068	-,134	,073	-,277	,010
	Equal variances not assumed			-1,840	212,995	,067	-,134	,073	-,277	,010

### Gesamtanzahl betroffener Frauen

#### Count

		Wochenbettkomplikationen		Total
		Nein	Ja	
code	Mimü	59	47	106
	Vomü	48	62	110
Total		107	109	216

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,122 <sup>a</sup>	1	,077	,102	,051	
Continuity Correction <sup>b</sup>	2,660	1	,103			
Likelihood Ratio	3,130	1	,077	,102	,051	
Fisher's Exact Test				,102	,051	
Linear-by-Linear Association	3,108 <sup>c</sup>	1	,078	,102	,051	,023
N of Valid Cases	216					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 52,51.

c. The standardized statistic is 1,763.

## Puberale Infektion

Count

		Puberale Infektion		Total
		Nein	Ja	
code	Mimü	105	1	106
	Vomü	108	2	110
Total		213	3	216

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,302 <sup>b</sup>	1	,583	1,000	,514	
Continuity Correction <sup>a</sup>	,000	1	1,000			
Likelihood Ratio	,308	1	,579	1,000	,514	
Fisher's Exact Test				1,000	,514	
Linear-by-Linear Association	,300 <sup>c</sup>	1	,584	1,000	,514	,384
N of Valid Cases	216					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,47.

c. The standardized statistic is ,548.

## Wundheilungsstörungen

Count

		Wundheilungsstörungen		Total
		Nein	Ja	
code	Mimü	106	0	106
	Vomü	109	1	110
Total		215	1	216

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,968 <sup>a</sup>	1	,325	1,000	,509	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,354	1	,245	1,000	,509	
Fisher's Exact Test				1,000	,509	
Linear-by-Linear Association	,964 <sup>c</sup>	1	,326	1,000	,509	,509
N of Valid Cases	216					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,49.

c. The standardized statistic is ,982.

## Mastitis

Count

		Mastitis		Total
		Nein	Ja	
code	Mimü	106	0	106
	Vomü	109	1	110
Total		215	1	216

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,968 <sup>b</sup>	1	,325	1,000	,509	
Continuity Correction <sup>a</sup>	,000	1	1,000			
Likelihood Ratio	1,354	1	,245	1,000	,509	
Fisher's Exact Test				1,000	,509	
Linear-by-Linear Association	,964 <sup>c</sup>	1	,326	1,000	,509	,509
N of Valid Cases	216					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,49.

c. The standardized statistic is ,982.

## Anämie

Count

		Anämie		Total
		Nein	Ja	
code	Mimü	59	47	106
	Vomü	49	61	110
Total		108	108	216

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,668 <sup>a</sup>	1	,102	,134	,067	
Continuity Correction <sup>b</sup>	2,242	1	,134			
Likelihood Ratio	2,673	1	,102	,134	,067	
Fisher's Exact Test				,134	,067	
Linear-by-Linear Association	2,655 <sup>c</sup>	1	,103	,134	,067	,029
N of Valid Cases	216					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 53,00.

c. The standardized statistic is 1,629.

weitere erfasste Komplikationen :

Count

		Toxisches Schocksyndrom	
		Nein	Total
code	Mimü	106	106
	Vomü	110	110
Total		216	216

Count

		Wochenbettdepression	
		Nein	Total
code	Mimü	106	106
	Vomü	110	110
Total		216	216

Count

		Thrombose	
		Nein	Total
code	Mimü	105	105
	Vomü	110	110
Total		215	215

Count

		Embolie	
		Nein	Total
code	Mimü	106	106
	Vomü	110	110
Total		216	216



### A3.4 Kindliches Outcome

#### Kindbezogene Komplikationen unter der Geburt

Kindliche Komplikationen gesamt

	code	N	Mean	Std. Deviation	Std. Error Mean
Kindliche Komplikationen	Mimü	110	1,98	1,459	,139
	Vomü	110	2,31	2,053	,196

#### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Kindliche Komplikationen	Equal variances assumed	6,215	,013	-1,363	218	,174	-,327	,240	-,801	,146
	Equal variances not assumed			-1,363	196,702	,174	-,327	,240	-,801	,146

Gesamtanzahl der Kinder mit Komplikationen

Count

		Gesamtanzahl der Kinder mit Komplikationen		Total
		Nein	Ja	
code	Mimü	36	72	108
	Vomü	39	71	110
Total		75	143	218

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,109 <sup>a</sup>	1	,742	,777	,426	
Continuity Correction <sup>b</sup>	,035	1	,852			
Likelihood Ratio	,109	1	,742	,777	,426	
Fisher's Exact Test				,777	,426	
Linear-by-Linear Association	,108 <sup>c</sup>	1	,742	,777	,426	,107
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 37,16.

c. The standardized statistic is -,329.

## abnorme Herzfrequenz

Count

		abnorme Herzfrequenz		Total
		Nein	Ja	
code	Mimü	69	39	108
	Vomü	79	31	110
Total		148	70	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,572 <sup>a</sup>	1	,210	,246	,134	
Continuity Correction <sup>b</sup>	1,229	1	,268			
Likelihood Ratio	1,574	1	,210	,246	,134	
Fisher's Exact Test				,246	,134	
Linear-by-Linear Association	1,565 <sup>c</sup>	1	,211	,246	,134	,053
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 34,68.

c. The standardized statistic is -1,251.

## Nabelschnurumschlingungen

Count

		Nabelschnurumschlingung		Total
		Nein	Ja	
code	Mimü	62	46	108
	Vomü	66	44	110
Total		128	90	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,151 <sup>d</sup>	1	,697	,783	,401	
Continuity Correction <sup>e</sup>	,063	1	,802			
Likelihood Ratio	,151	1	,697	,783	,401	
Fisher's Exact Test				,783	,401	
Linear-by-Linear Association	,150 <sup>c</sup>	1	,698	,783	,401	,101
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 44,59.

c. The standardized statistic is -,388.

## Insertio velamentosa

## Count

		Insertio velamentosa		Total
		Nein	Ja	
code	Mimü	108	0	108
	Vomü	108	2	110
Total		216	2	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,982 <sup>d</sup>	1	,159	,498	,253	
Continuity Correction <sup>e</sup>	,486	1	,486			
Likelihood Ratio	2,754	1	,097	,498	,253	
Fisher's Exact Test				,498	,253	
Linear-by-Linear Association	1,973 <sup>c</sup>	1	,160	,498	,253	,253
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,99.

c. The standardized statistic is 1,405.

## Count

		Kephalhämatom		Total
		Nein	Ja	
code	Mimü	106	2	108
	Vomü	108	2	110
Total		214	4	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,000 <sup>u</sup>	1	,985	1,000	,682	
Continuity Correction <sup>ß</sup>	,000	1	1,000			
Likelihood Ratio	,000	1	,985	1,000	,682	
Fisher's Exact Test				1,000	,682	
Linear-by-Linear Association	,000 <sup>c</sup>	1	,985	1,000	,682	,378
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,98.

c. The standardized statistic is -,018.

## Caput succedaneum

## Count

		caput succedaneum		Total
		Nein	Ja	
code	Mimü	104	4	108
	Vomü	110	0	110
Total		214	4	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,150 <sup>p</sup>	1	,042	,059	,059	
Continuity Correction <sup>ß</sup>	2,349	1	,125			
Likelihood Ratio	5,695	1	,017	,059	,059	
Fisher's Exact Test				,059	,059	
Linear-by-Linear Association	4,131 <sup>c</sup>	1	,042	,059	,059	,059
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,98.

c. The standardized statistic is -2,033.

## obere Plexuslähmung

Count

		obere Plexuslähmung	
		Nein	Total
code	Mimü	108	108
	Vomü	110	110
Total		218	218

untere Plexuslähmung

Count

		untere Plexuslähmung	
		Nein	Total
code	Mimü	108	108
	Vomü	110	110
Total		218	218

Klavikularfraktur

Count

		Klavikularfraktur		Total
		Nein	Ja	
code	Mimü	108	0	108
	Vomü	109	1	110
Total		217	1	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,986 <sup>a</sup>	1	,321	1,000	,505	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,373	1	,241	1,000	,505	
Fisher's Exact Test				1,000	,505	
Linear-by-Linear Association	,982 <sup>c</sup>	1	,322	1,000	,505	,505
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is ,991.

Konjungtivitis

---

Count

		Konjungtivitis	
		Nein	Total
code	Mimü	108	108
	Vomü	110	110
Total		218	218

## andere Komplikationen

Count

		andere Komplikationen		Total
		Nein	Ja	
code	Mimü	105	3	108
	Vomü	93	17	110
Total		198	20	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	10,510 <sup>a</sup>	1	,001	,002	,001	
Continuity Correction <sup>b</sup>	9,044	1	,003			
Likelihood Ratio	11,527	1	,001	,002	,001	
Fisher's Exact Test				,002	,001	
Linear-by-Linear Association	10,462 <sup>c</sup>	1	,001	,002	,001	,001
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9,91.

c. The standardized statistic is 3,234.

## PH-Wert des Kindes bei der Geburt

## PH

code			Frequency	Percent	Valid Percent	Cumulative Percent
Mimü	Valid	> 7,3	47	42,3	45,2	45,2
		7,2 - 7,3	52	46,8	50,0	95,2
		7,1 - 7,2	5	4,5	4,8	100,0
		Total	104	93,7	100,0	
	Missing	99	7	6,3		
Total			111	100,0		
Vomü	Valid	> 7,3	53	48,2	48,2	48,2
		7,2 - 7,3	55	50,0	50,0	98,2
		7,1 - 7,2	2	1,8	1,8	100,0
		Total	110	100,0	100,0	

Count

		leichte Azidose		Total
		Nein	Ja	
code	Mimü	99	5	104
	Vomü	108	2	110
Total		207	7	214

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,510 <sup>a</sup>	1	,219	,269	,200	
Continuity Correction <sup>b</sup>	,713	1	,398			
Likelihood Ratio	1,551	1	,213	,269	,200	
Fisher's Exact Test				,269	,200	
Linear-by-Linear Association	1,503 <sup>c</sup>	1	,220	,269	,200	,149
N of Valid Cases	214					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 3,40.

c. The standardized statistic is -1,226.

## Base-Excess

## BE

code			Frequency	Percent	Valid Percent	Cumulative Percent
Mimü	Valid	+5 - (-5)	86	77,5	82,7	82,7
		-5 - (-10)	18	16,2	17,3	100,0
		Total	104	93,7	100,0	
Missing	Total	99	7	6,3		
			111	100,0		
Vomü	Valid	+5 - (-5)	100	90,9	93,5	93,5
		-5 - (-10)	7	6,4	6,5	100,0
		Total	107	97,3	100,0	
Missing	Total	99	3	2,7		
			110	100,0		



## Körperlicher Entwicklungsstand des Neugeborenen

Geburtsgewicht:

gewicht

Mimü	N	Valid	110
		Missing	1
	Mean		3183,32
	Median		3280,00
	Std. Deviation		714,445
	Minimum		745
	Maximum		5000
Vomü	N	Valid	110
		Missing	0
	Mean		3231,91
	Median		3250,00
	Std. Deviation		689,402
	Minimum		1180
	Maximum		4620

niedriges Geburtsgewicht

	code	N	Mean	Std. Deviation	Std. Error Mean
LBW	Gewicht	Mimü	18	2076,39	104,662
		Vomü	13	1841,54	129,628
LBW	Gewicht	Mimü	92	3399,89	55,636
		Vomü	97	3418,25	47,131

### Independent Samples Test

			Levene's Test for Equality of Variances		t-test for Equality of Means						
			F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
										Lower	Upper
LBW	Gewicht	Equal variances assumed	,660	,423	1,422	29	,166	234,850	165,188	-102,998	572,699
		Equal variances not assumed			1,410	25,189	,171	234,850	166,605	-108,149	577,850
non LBW	Gewicht	Equal variances assumed	,311	,578	-,253	187	,801	-18,356	72,648	-161,670	124,958
		Equal variances not assumed			-,252	180,404	,802	-18,356	72,916	-162,234	125,521

## sehr niedriges Geburtsgewicht

code			N	Mean	Std. Deviation	Std. Error Mean
VLBW	Gewicht	Mimü	2	1112,50	519,723	367,500
		Vomü	4	1240,00	76,920	38,460
non VLBW	Gewicht	Mimü	108	3221,67	660,158	63,524
		Vomü	106	3307,08	579,963	56,331

## Independent Samples Test

			Levene's Test for Equality of Variances		t-test for Equality of Means						
			F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
VLBWcode										Lower	Upper
VLBW	Gewicht	Equal variances assumed	113,267	,000	-,549	4	,612	-127,500	232,324	-772,534	517,534
		Equal variances not assumed			-,345	1,022	,788	-127,500	369,507	-4590,471	4335,471
non VLBW	Gewicht	Equal variances assumed	1,855	,175	-1,005	212	,316	-85,409	85,005	-252,973	82,155
		Equal variances not assumed			-1,006	209,458	,316	-85,409	84,903	-252,782	81,964

## Geburtsgröße

## Größe

Mimü	N	Valid	110
		Missing	1
	Mean		49,05
	Median		49,50
	Std. Deviation		3,037
	Minimum		33
	Maximum		55
Vomü	N	Valid	110
		Missing	0
	Mean		48,94
	Median		49,00
	Std. Deviation		3,713
	Minimum		29
	Maximum		54

## Kopfumfang

Kopfumfang			
Mimü	N	Valid	107
		Missing	4
	Mean		34,320
	Median		34,500
	Std. Deviation		2,6154
	Minimum		24,5
	Maximum		51,0
Vomü	N	Valid	110
		Missing	0
	Mean		34,191
	Median		34,000
	Std. Deviation		2,4438
	Minimum		24,0
	Maximum		45,0

## Reifestatus des Neugeborenen

## Hypotrophe Neugeborene

Count				
		hypotrophe Neugeborene		Total
		Nein	Ja	
code	Mimü	90	11	101
	Vomü	98	12	110
Total		188	23	211

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,000 <sup>b</sup>	1	,997	1,000	,587	
Continuity Correction <sup>a</sup>	,000	1	1,000			
Likelihood Ratio	,000	1	,997	1,000	,587	
Fisher's Exact Test				1,000	,587	
Linear-by-Linear Association	,000 <sup>c</sup>	1	,997	1,000	,587	,174
N of Valid Cases	211					

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11,01.

c. The standardized statistic is ,004.

## eutrophe Neugeborenen

Count

		eutrophe Neugeborenen		Total
		Nein	Ja	
code	Mimü	22	79	101
	Vomü	39	71	110
Total		61	150	211

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,789 <sup>a</sup>	1	,029	,034	,020	
Continuity Correction <sup>b</sup>	4,147	1	,042			
Likelihood Ratio	4,844	1	,028	,034	,020	
Fisher's Exact Test				,034	,020	
Linear-by-Linear Association	4,767 <sup>c</sup>	1	,029	,034	,020	,011
N of Valid Cases	211					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 29,20.

c. The standardized statistic is -2,183.

## Hypertrophe Neugeborene

Count

		Hypertrophe Neugeborene		Total
		Nein	Ja	
code	Mimü	95	6	101
	Vomü	92	18	110
Total		187	24	211

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	5,675 <sup>a</sup>	1	,017	,028	,014	
Continuity Correction <sup>b</sup>	4,688	1	,030			
Likelihood Ratio	5,943	1	,015	,018	,014	
Fisher's Exact Test				,018	,014	
Linear-by-Linear Association	5,648 <sup>c</sup>	1	,017	,028	,014	,010
N of Valid Cases	211					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 11,49.

c. The standardized statistic is 2,376.

## Hypotrophe Frühgeborene

Count

		Hypotrophe Früchen		Total
		Nein	Ja	
code	Mimü	10	4	14
	Vomü	9	1	10
Total		19	5	24

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,220 <sup>a</sup>	1	,269	,358	,283	
Continuity Correction <sup>b</sup>	,354	1	,552			
Likelihood Ratio	1,310	1	,252	,358	,283	
Fisher's Exact Test				,358	,283	
Linear-by-Linear Association	1,169 <sup>c</sup>	1	,280	,358	,283	,236
N of Valid Cases	24					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,08.

c. The standardized statistic is -1,081.

## Eutrophe Frühgeborene

Count

		eutrophe Früchen		Total
		Nein	Ja	
code	Mimü	5	9	14
	Vomü	3	7	10
Total		8	16	24

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,086 <sup>a</sup>	1	,770	1,000	,561	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,086	1	,769	1,000	,561	
Fisher's Exact Test				1,000	,561	
Linear-by-Linear Association	,082 <sup>c</sup>	1	,774	1,000	,561	,327
N of Valid Cases	24					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 3,33.

c. The standardized statistic is ,287.

## Hypertrophe Frühgeborene

## Crosstab

Count

		Hypertrophe Früchen		Total
		Nein	Ja	
code	Mimü	14	0	14
	Vomü	8	2	10
Total		22	2	24

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,055 <sup>a</sup>	1	,081	,163	,163	
Continuity Correction <sup>b</sup>	,997	1	,318			
Likelihood Ratio	3,760	1	,052	,163	,163	
Fisher's Exact Test				,163	,163	
Linear-by-Linear Association	2,927 <sup>c</sup>	1	,087	,163	,163	,163
N of Valid Cases	24					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,83.

c. The standardized statistic is 1,711.

## Überreifezeichen

		Überreifezeichen		Total
		Nein	Ja	
code	Mimü	54	29	83
	Vomü	77	15	92
Total		131	44	175

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	8,051 <sup>a</sup>	1	,005	,005	,004	
Continuity Correction <sup>b</sup>	7,091	1	,008			
Likelihood Ratio	8,129	1	,004	,005	,004	
Fisher's Exact Test				,005	,004	
Linear-by-Linear Association	8,005 <sup>c</sup>	1	,005	,005	,004	,003
N of Valid Cases	175					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 20,87.

c. The standardized statistic is -2,829.

## APGAR

code			Apgar nach 1 Minute	Apgar nach 5 Minuten	Apgar nach 10 Minuten
Mimü	N	Valid	109	109	109
		Missing	2	2	2
	Mean	8,47	9,47	9,78	
	Median	9,00	10,00	10,00	
	Std. Deviation	1,337	,800	,567	
	Minimum	1	6	7	
	Maximum	10	10	10	
Vomü	N	Valid	110	110	110
		Missing	0	0	0
	Mean	8,45	9,43	9,74	
	Median	9,00	10,00	10,00	
	Std. Deviation	,982	,772	,536	
	Minimum	3	6	8	
	Maximum	9	10	10	

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Apgar nach 1 Minute	Equal variances assumed	,476	,491	,142	217	,888	,022	,158	-,290	,335
	Equal variances not assumed			,141	198,216	,888	,022	,159	-,290	,335
Apgar nach 5 Minuten	Equal variances assumed	,023	,881	,382	217	,703	,041	,106	-,169	,250
	Equal variances not assumed			,382	216,557	,703	,041	,106	-,169	,250
Apgar nach 10 Minuten	Equal variances assumed	,723	,396	,583	217	,561	,043	,075	-,104	,190
	Equal variances not assumed			,583	216,097	,561	,043	,075	-,104	,190

## Notfallmaßnahmen

### O<sup>2</sup>- Gabe unter der Geburt

Count

		O2- Gabe		Total
		Nein	Ja	
code	Mimü	94	15	109
	Vomü	97	12	109
Total		191	27	218

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,380 <sup>a</sup>	1	,537	,682	,341	
Continuity Correction <sup>b</sup>	,169	1	,681			
Likelihood Ratio	,381	1	,537	,682	,341	
Fisher's Exact Test				,682	,341	
Linear-by-Linear Association	,379 <sup>c</sup>	1	,538	,682	,341	,135
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 13,50.

c. The standardized statistic is -,615.



## Medikamentöse Soforttherapie

Count		Medikamente		Total
		Nein	Ja	
code	Mimü	104	5	109
	Vomü	98	11	109
Total		202	16	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,428 <sup>a</sup>	1	,119	,193	,096	
Continuity Correction <sup>b</sup>	1,686	1	,194			
Likelihood Ratio	2,484	1	,115	,193	,096	
Fisher's Exact Test				,193	,096	
Linear-by-Linear Association	2,417 <sup>c</sup>	1	,120	,193	,096	,063
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 8,00.

c. The standardized statistic is 1,555.

## Reanimation unter der Geburt

Count		Reanimation		Total
		Nein	Ja	
code	Mimü	107	2	109
	Vomü	109	0	109
Total		216	2	218

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,019 <sup>a</sup>	1	,155	,498	,249	
Continuity Correction <sup>b</sup>	,505	1	,477			
Likelihood Ratio	2,791	1	,095	,498	,249	
Fisher's Exact Test				,498	,249	
Linear-by-Linear Association	2,009 <sup>c</sup>	1	,156	,498	,249	,249
N of Valid Cases	218					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is -1,417.

## Lungenreife beim Kind durchgeführt

Count

		Lungenreife Kind		Total
		Nein	Ja	
code	Mimü	111	0	111
	Vomü	106	4	110
Total		217	4	221

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3,175 <sup>a</sup>	1	,075	,125	,108	
Continuity Correction <sup>b</sup>	1,500	1	,221			
Likelihood Ratio	4,586	1	,032	,125	,108	
Fisher's Exact Test				,125	,108	
Linear-by-Linear Association	3,048 <sup>c</sup>	1	,081	,125	,108	,108
N of Valid Cases	25					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,60.

c. The standardized statistic is 1,746.

## Aufenthalt auf der neonatologischen Station

### Wie lange auf neonatologischen Station

code			Frequency	Percent	Valid Percent	Cumulative Percent
Mimü	Valid	0	1	,9	3,6	3,6
		unter 3 Tagen	5	4,5	17,9	21,4
		3-5 Tage	3	2,7	10,7	32,1
		5-10 Tage	11	9,9	39,3	71,4
		über 10 Tage	8	7,2	28,6	100,0
	Missing	Total	28	25,2	100,0	
		88	79	71,2		
		unbekannt	3	2,7		
		System	1	,9		
		Total	83	74,8		
Total		111	100,0			
vomü	Valid	unter 3 Tagen	5	4,5	20,8	20,8
		3-5 Tage	2	1,8	8,3	29,2
		5-10 Tage	7	6,4	29,2	58,3
		über 10 Tage	10	9,1	41,7	100,0
		Total	24	21,8	100,0	
	Missing	88	84	76,4		
		unbekannt	2	1,8		
		Total	86	78,2		
Total		110	100,0			

## Neonatologischer Aufenthalt

Count

		Neonatologie		Total
		Nein	Ja	
code	Mimü	82	26	108
	Vomü	83	26	109
Total		165	52	217

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,001 <sup>d</sup>	1	,970	1,000	,548	
Continuity Correction <sup>e</sup>	,000	1	1,000			
Likelihood Ratio	,001	1	,970	1,000	,548	
Fisher's Exact Test				1,000	,548	
Linear-by-Linear Association	,001 <sup>c</sup>	1	,970	1,000	,548	,126
N of Valid Cases	217					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 25,88.

c. The standardized statistic is -,038.

## Entlassung gegen ärztlichen Rat

## Entlassung gegen ärztlichen Rat

code			Frequency	Percent	Valid Percent	Cumulative Percent
Mimü	Valid	Nein	102	91,9	95,3	95,3
		Ja	5	4,5	4,7	100,0
		Total	107	96,4	100,0	
	Missing	unbekannt	4	3,6		
Total			111	100,0		
Vomü	Valid	Nein	104	94,5	95,4	95,4
		Ja	5	4,5	4,6	100,0
		Total	109	99,1	100,0	
	Missing	unbekannt	1	,9		
	Total			110	100,0	

Count

		Entlassung gegen ärztlichen Rat		Total
		Nein	Ja	
code	Mimü	102	5	107
	Vomü	104	5	109
Total		206	10	216

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,001 <sup>d</sup>	1	,976	1,000	,614	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,001	1	,976	1,000	,614	
Fisher's Exact Test				1,000	,614	
Linear-by-Linear Association	,001 <sup>c</sup>	1	,976	1,000	,614	,252
N of Valid Cases	216					

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,95.

c. The standardized statistic is -,030.

## Untersuchungsergebnisse des Neugeborenen

## Hüftsonographie

Count

		Risiko Hüftsono		Total
		kein Risiko	Risiko	
code	Mimü	57	30	87
	Vomü	74	26	100
Total		131	56	187

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,596 <sup>a</sup>	1	,207	,263	,135	
Continuity Correction <sup>b</sup>	1,217	1	,270			
Likelihood Ratio	1,594	1	,207	,263	,135	
Fisher's Exact Test				,263	,135	
Linear-by-Linear Association	1,587 <sup>c</sup>	1	,208	,263	,135	,058
N of Valid Cases	187					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 26,05.

c. The standardized statistic is -1,260.

## Schädelsonographie

## Count

		Risiko Schädelsono		Total
		kein Risiko	Risiko	
code	Mimü	78	9	87
	Vomü	93	8	101
Total		171	17	188

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,334 <sup>b</sup>	1	,563	,616	,372	
Continuity Correction <sup>b</sup>	,104	1	,747			
Likelihood Ratio	,333	1	,564	,616	,372	
Fisher's Exact Test				,616	,372	
Linear-by-Linear Association	,332 <sup>c</sup>	1	,564	,616	,372	,170
N of Valid Cases	188					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,87.

c. The standardized statistic is -,576.

## Neugeborenen-screening

Count

		Risiko Neugeborenen-screening		Total
		kein Risiko	Risiko	
code	Mimü	45	2	47
	Vomü	40	2	42
Total		85	4	89

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,013 <sup>a</sup>	1	,908	1,000	,648	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,013	1	,908	1,000	,648	
Fisher's Exact Test				1,000	,648	
Linear-by-Linear Association	,013 <sup>c</sup>	1	,909	1,000	,648	,381
N of Valid Cases	89					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,89.

c. The standardized statistic is ,115.

## Hörscreening

Count

		Risiko Hörscreening		Total
		kein Risiko	Risiko	
code	Mimü	72	1	73
	Vomü	74	4	78
Total		146	5	151

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,664 <sup>d</sup>	1	,197	,368	,205	
Continuity Correction <sup>e</sup>	,697	1	,404			
Likelihood Ratio	1,789	1	,181	,368	,205	
Fisher's Exact Test				,368	,205	
Linear-by-Linear Association	1,653 <sup>c</sup>	1	,199	,368	,205	,170
N of Valid Cases	151					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,42.

c. The standardized statistic is 1,286.

## Weitere pathologische Befunde des Neugeborenen

veränderte Bilirubinwerte

Count

		Ikterus des Kindes		Total
		Nein	Ja	
code	Mimü	91	18	109
	Vomü	99	11	110
Total		190	29	219

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2,022 <sup>d</sup>	1	,155	,168	,111	
Continuity Correction <sup>e</sup>	1,495	1	,221			
Likelihood Ratio	2,039	1	,153	,168	,111	
Fisher's Exact Test				,168	,111	
Linear-by-Linear Association	2,013 <sup>c</sup>	1	,156	,168	,111	,059
N of Valid Cases	219					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,43.

c. The standardized statistic is -1,419.



## pathologische Laborwerte des Kindes gesamt

	code	N	Mean	Std. Deviation	Std. Error Mean
Laborwerte	Mimü	111	,04	,187	,018
Kind	Vomü	110	,24	,468	,045

## Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Laborwerte	Equal variances assumed	89,698	,000	-4,186	219	,000	-,200	,048	-,295	-,106
Kind	Equal variances not assumed			-4,172	142,784	,000	-,200	,048	-,295	-,105

## Gesamtanzahl der Kinder mit pathologischen Laborwerten

## Count

	code	Laborwerte des Kindes Gesamtveränderung		Total
		Nein	Ja	
	Mimü	106	4	110
	Vomü	86	24	110
Total		192	28	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	16,369 <sup>a</sup>	1	,000	,000	,000	
Continuity Correction <sup>b</sup>	14,773	1	,000			
Likelihood Ratio	17,937	1	,000	,000	,000	
Fisher's Exact Test				,000	,000	
Linear-by-Linear Association	16,295 <sup>c</sup>	1	,000	,000	,000	,000
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 14,00.

c. The standardized statistic is 4,037.

## pathologische Eisenwerte

Count

		Kindlicher Eisenwert		Total
		Nein	Ja	
code	Mimü	109	1	110
	Vomü	110	0	110
Total		219	1	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1,005 <sup>a</sup>	1	,316	1,000	,500	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	1,391	1	,238	1,000	,500	
Fisher's Exact Test				1,000	,500	
Linear-by-Linear Association	1,000 <sup>c</sup>	1	,317	1,000	,500	,500
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,50.

c. The standardized statistic is -1,000.

## pathologische Gerinnungswerte

Count

		Gerinnungswerte des Kind		Total
		Nein	Ja	
code	Mimü	109	1	110
	Vomü	109	1	110
Total		218	2	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,000 <sup>a</sup>	1	1,000	1,000	,751	
Continuity Correction <sup>b</sup>	,000	1	1,000			
Likelihood Ratio	,000	1	1,000	1,000	,751	
Fisher's Exact Test				1,000	,751	
Linear-by-Linear Association	,000 <sup>c</sup>	1	1,000	1,000	,751	,502
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

c. The standardized statistic is ,000.

## pathologische Glucosewerte

Count

		Glucosewerte des Kindes		Total
		Nein	Ja	
code	Mimü	110	0	110
	Vomü	100	10	110
Total		210	10	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	10,476 <sup>a</sup>	1	,001	,002	,001	
Continuity Correction <sup>b</sup>	8,486	1	,004			
Likelihood Ratio	14,339	1	,000	,002	,001	
Fisher's Exact Test				,002	,001	
Linear-by-Linear Association	10,429 <sup>c</sup>	1	,001	,002	,001	,001
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 5,00.

c. The standardized statistic is 3,229.

## pathologische Blutwerte

Count

		andere Blutwerte des Kindes		Total
		Nein	Ja	
code	Mimü	108	2	110
	Vomü	95	15	110
Total		203	17	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	10,774 <sup>a</sup>	1	,001	,002	,001	
Continuity Correction <sup>b</sup>	9,180	1	,002			
Likelihood Ratio	12,085	1	,001	,002	,001	
Fisher's Exact Test				,002	,001	
Linear-by-Linear Association	10,725 <sup>c</sup>	1	,001	,002	,001	,001
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 8,50.

c. The standardized statistic is 3,275.

weitere pathologische Befunde des Neugeborenen.

Count

		weitere Pathologien		Total
		Nein	Ja	
code	Mimü	90	21	111
	Vomü	84	26	110
Total		174	47	221

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	,734 <sup>d</sup>	1	,391	,415	,244	
Continuity Correction <sup>e</sup>	,480	1	,489			
Likelihood Ratio	,735	1	,391	,415	,244	
Fisher's Exact Test				,415	,244	
Linear-by-Linear Association	,731 <sup>c</sup>	1	,393	,415	,244	,091
N of Valid Cases	221					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 23,39.

c. The standardized statistic is ,855.

### A3.5 Jugendamt oder Vormundschaft

Count

		Jugendamt Informiert		Total
		Nein	Ja	
code	Mimü	93	17	110
	Vomü	106	4	110
Total		199	21	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	8,897 <sup>u</sup>	1	,003	,005	,002	
Continuity Correction <sup>ß</sup>	7,581	1	,006			
Likelihood Ratio	9,512	1	,002	,005	,002	
Fisher's Exact Test				,005	,002	
Linear-by-Linear Association	8,856 <sup>c</sup>	1	,003	,005	,002	,002
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 10,50.

c. The standardized statistic is -2,976.

## Count

		Vormund		Total
		Nein	Ja	
code	Mimü	106	4	110
	Vom ü	110	0	110
Total		216	4	220

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	4,074 <sup>u</sup>	1	,044	,122	,061	
Continuity Correction <sup>ß</sup>	2,292	1	,130			
Likelihood Ratio	5,619	1	,018	,122	,061	
Fisher's Exact Test				,122	,061	
Linear-by-Linear Association	4,056 <sup>c</sup>	1	,044	,122	,061	,061
N of Valid Cases	220					

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,00.

c. The standardized statistic is -2,014.

### A3.6 Diskussion

	code	N	Mean	Std. Deviation	Std. Error Mean
Compositescore	Mimü	108	1,87	1,368	,132
Kindliches Outcome	Vomü	110	2,13	1,926	,184
Compositescore	Mimü	109	,91	1,295	,124
Erkrankungen in der SS	Vomü	109	1,17	1,779	,170
Compositescore	Mimü	105	2,46	,920	,090
Geburtskomplikationen	Vomü	110	3,07	1,055	,101

#### Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Compositescore Kindliches Outcome	Equal variances assumed	5,849	,016	-1,134	216	,258	-,257	,227	-,704	,190
	Equal variances not assumed			-1,137	196,850	,257	-,257	,226	-,702	,189
Compositescore Erkrankungen in der SS	Equal variances assumed	6,828	,010	-1,263	216	,208	-,266	,211	-,681	,149
	Equal variances not assumed			-1,263	197,371	,208	-,266	,211	-,682	,150
Compositescore Geburtskomplikationen	Equal variances assumed	,766	,383	-4,550	213	,000	-,616	,135	-,882	-,349
	Equal variances not assumed			-4,564	211,290	,000	-,616	,135	-,881	-,350

#### Größe der Frauen zur Geburt

##### Group Statistics

	code	N	Mean	Std. Deviation	Std. Error Mean
Größe der Mutter	Mimü	110	166,818	7,0713	,6742
	Vomü	110	167,873	6,3247	,6030

## Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Größe der Mutter	Equal variances assumed	,506	,478	-1,166	218	,245	-1,0545	,9046	-2,8373	,7283
	Equal variances not assumed			-1,166	215,341	,245	-1,0545	,9046	-2,8375	,7284