

**Nitrogen and organic matter cycling in coastal systems: case studies from the  
Baltic Sea**

Dissertation  
zur  
Erlangung des akademischen Grades  
Doktor der Naturwissenschaften  
(doctor rerum naturalium)

vorgelegt der  
Mathematisch- Naturwissenschaftlichen Fakultät  
der Universität Rostock

von  
Frederike Korth,  
geb. am 28.06.1982 in Preetz

Rostock 2012

urn:nbn:de:gbv:28-diss2013-0003-7

Die vorliegende Arbeit wurde von der Mathematisch-Naturwissenschaftlichen Fakultät der Universität Rostock als Dissertation angenommen und am 22. Oktober 2012 erfolgreich verteidigt.

Die Gutachter waren:

PD Dr. Maren Voß, Leibniz-Institut für Ostseeforschung Warnemünde an der Universität Rostock

Prof. Dr. Christoph Humborg, Stockholm University, Sweden

**Contents**

<b>Summary</b> .....	III
<b>Zusammenfassung</b> .....	V
<b>1.Introduction</b> .....	1
1.1 Ecological problems of coastal eutrophication.....	1
1.2 Global nitrogen cycle.....	2
1.3 Dissolved organic matter as part of the nitrogen cycle.....	5
1.4 Stable isotopes as a tool to unravel the nitrogen cycle.....	7
1.4.1 Tracking nitrogen in coastal areas .....	8
1.5 The Baltic Sea and its sensitivity towards eutrophication.....	11
1.6 Aim of the study.....	13
<b>2.History and scenarios of future development of Baltic Sea eutrophication</b> .....	14
<b>3.Nitrogen turnover during spring outflow from the nitrate-rich Curonian and Szczecin lagoon using dual isotopes in nitrate</b> .....	15
<b>4.Uptake of dissolved organic nitrogen by size-fractionated plankton along a salinity gradient from the North Sea to the Baltic Sea</b> .....	16
<b>5.Tracing inputs of terrestrial high molecular weight dissolved organic matter within the Baltic Sea Ecosystem</b> .....	17
<b>6.Conclusions and Perspectives</b> .....	18
<b>Bibliography</b> .....	23

## Summary

The depletion of dissolved oxygen in bottom waters and the expansion of harmful algae blooms (HAB's) are most common responses of aquatic ecosystems to eutrophication, and thus also to the semi-enclosed, brackish Baltic Sea with high riverine nutrient loads. In the first part of this thesis the historical progression of eutrophication in the Baltic Sea, and the reasons for its sensitivity towards eutrophication are summarized. The northern watersheds, which are sparsely populated and mainly covered by boreal forests, were compared to the southern watersheds, which are surrounded by highly industrialized and densely populated areas that are mainly in agricultural use. The evaluation of long term data sets and model results indicates that in addition to changes in nutrient inputs, increased temperature and precipitation are likely to become important forcings for the Baltic Sea. Moreover, it has been suggested that lagoons and near shore areas remove large quantities of riverine nitrogen, but will only be able to do so as long as they remain oxic meaning that this system service is threatened due to increasing coastal hypoxia. In the second study the nitrogen removal processes in the adjacent coastal areas of the Oder and Nemunas rivers were characterized during peak outflow, by means of dual stable isotopes measurements in nitrate ( $\delta^{15}\text{N}\text{-NO}_3^-$  and  $\delta^{18}\text{O}\text{-NO}_3^-$ ) and nitrate uptake rates, since nitrate is highly bioavailable and the dominant nitrogen component delivered by these rivers. Results show that the isotopic signal of nitrate is dominated on the one hand by mixing and assimilation in the surface waters and on the other hand by denitrification in the near bottom waters. Calculated fractionation factors ( $^{15}\epsilon$  and  $^{18}\epsilon$ ) of around 10‰ in the near bottom waters infer that the isotopic enrichment from sedimentary denitrification may be higher in permeable, sandy than in muddy sediments. So far, permeable sediments with low organic content were mainly considered to contribute little to the biogeochemical cycling. However, the isotope data of nitrate from this study indicate that denitrification in these sediments may be an important process. Because permeable sediments account for up to 70% of the continental shelf area their contribution to the Baltic Sea loss budget as well as to the global nitrogen loss budget should be reevaluated.

Large quantities of nitrogen are also delivered to the Baltic Sea in form of dissolved organic nitrogen (DON) via rivers and nitrogen fixation. However, the role of DON in the nitrogen cycle is still poorly understood and its impact on eutrophication has not been further considered in nitrogen load reductions. Therefore, in the third study the major aim was to evaluate the role of DON as a potential nutrient source for surface plankton in summer when dissolved inorganic nitrogen (DIN) concentrations are low. Two different  $^{15}\text{N}$  labeled bulk DON substrates were produced and DON uptake rates were determined along a salinity gradient from the North Sea to the Baltic Sea. Uptake rates in the Baltic Sea (184 to 1213 nmol N l<sup>-1</sup> h<sup>-1</sup>) were an order of magnitude higher compared to rates in the North Sea or the Chesapeake Bay (USA). The findings from this study indicate that DON is

an important component of plankton nutrition and can fuel primary production and therefore also contribute to eutrophication. Additionally, the conservative versus non-conservative behavior of dissolved organic matter (DOM) including DON and DOC (dissolved organic carbon) was tested along the same salinity gradient and the share of terrestrial DOM to the total DOM pool was determined by means of  $\delta^{13}\text{C}$  values in the fourth study. Results indicate that up to 83% of the DOM is derived from terrestrial sources in the Baltic Sea and that substantial amounts of DOM (>50%) delivered by rivers are degraded near the coastline. Since eutrophication is a global problem, results from this thesis can be translated to other coastal zones of temperate climate with similar nitrogen loads. Overall, it seems necessary to reevaluate the role of sediments for the nitrogen turnover and its nitrogen removal capacity and to consider dissolved organic substances as potential nutrient sources.

## Zusammenfassung

Die Verarmung an gelösten Sauerstoff im bodennahen Wasser und die Ausbreitung von toxischen Algenblüten sind die am häufigsten auftretenden Folgen von Eutrophierung in aquatischen Ökosystemen, so auch in der Ostsee, einem der größten Brackwassergebiete der Welt mit hohen Nährstofffrachten aus Flusseinträgen. Der erste Teil dieser Doktorarbeit umfasst eine Literaturstudie in der die geschichtliche Entwicklung der Eutrophierung und die Gründe warum die Ostsee besonders sensibel auf die vom Menschen verursachte Belastungen reagiert, zusammengefasst sind. Hierbei wurden die nördlichen Einzugsgebiete der Ostsee, die nur dünn besiedelt und hauptsächlich von Wäldern bedeckt sind, mit den südlichen Einzugsgebieten, die stark besiedelt und von Industrie und Landwirtschaft geprägt sind, verglichen. Die Auswertung von Langzeitdatenmessungen und von Modellierungsstudien ergab, dass wahrscheinlich neben sich ändernden Nährstoffeinträgen auch steigende Temperaturen und Niederschläge in Zukunft eine größere Rolle in der Ostsee spielen werden. Außerdem wurde vermutet, dass Haie, die normalerweise große Mengen an über Flüsse eingetragene Stickstofffrachten entfernen, dies nur so lange bewerkstelligen können wie die Küstengebiete oxisch bleiben. Durch die zunehmende Sauerstoffverarmung in Folge der Eutrophierung in Küstengebieten ist diese Funktion aber gefährdet. Im zweiten Teil dieser Arbeit wurden Prozesse die Stickstoff aus dem angrenzenden Küstengebieten der Flüsse Oder und Nemunas während der Hauptabflusszeit im Frühjahr entfernen, charakterisiert. Da Nitrat zu dieser Jahreszeit die Hauptstickstoffkomponente ist, erfolgte dieses mit Hilfe von Messungen an stabilen Isotopen im Nitrat ( $\delta^{15}\text{N-NO}_3^-$  and  $\delta^{18}\text{O-NO}_3^-$ ) und Nitrat Aufnahmeraten. Ergebnisse zeigen, dass das Isotopensignal von Nitrat auf der einen Seite im Oberflächenwasser durch Mischung und Nitrat Aufnahme dominiert ist und auf der anderen Seite im Bodennahem Wasser durch Denitrifizierung. Berechnete Fraktionierungsfaktoren ( $^{15}\epsilon$  and  $^{18}\epsilon$ ) von 10‰ im bodennahen Wasser lassen darauf schließen, dass die Anreicherung in den Isotopen durch Denitrifizierung in permeablen Sedimenten höher ist als in feinen Sedimenten. Bisher fanden permeable Sedimente mit niedrigem organischem Gehalt in biogeochemischen Kreisläufen wenig Beachtung, aber die Daten von den Isotopenmessungen lassen darauf schließen, dass auch diese Sedimente durch Denitrifizierung einen wichtigen Anteil zur Stickstoffentfernung beitragen. Da permeable Sedimente bis zu 70% der Kontinentalen Schelfmeere ausmachen, müsste ihr Beitrag sowohl in Stickstoff Budgets der Ostsee wie in globalen überdacht werden.

Stickstoff wird nicht nur in Form von gelöstem anorganischem Stickstoff (DIN) über Flüsse und Stickstoff Fixierung in die Ostsee eingetragen sondern auch in Form von gelöstem organischem Stickstoff (DON). Die Rolle von DON im Stickstoffkreislauf ist immer noch wenig verstanden und in Bezug auf Eutrophierung fand es bisher wenig Beachtung. Deshalb wurde in der dritten Studie dieser Doktorarbeit die Rolle des DON im Sommer, wenn alle anorganischen Stickstoffquellen verbraucht

sind, als potentielle Nährstoffquelle für das Plankton im Oberflächenwasser untersucht. Zwei unterschiedliche  $^{15}\text{N}$  markierte DON Substrate wurden produziert und Aufnahmeraten von diesen beiden Tracern entlang eines Salinitätsgradienten in der Nord- und Ostsee bestimmt. Die Aufnahmeraten ( $184 \text{ to } 1213 \text{ nmol N l}^{-1} \text{ h}^{-1}$ ) in der Ostsee waren doppelt so hoch wie die in der Nordsee oder auch im Vergleich zur Chesapeake Bay (USA). Die Ergebnisse dieser Studie lassen darauf schließen, dass DON eine wichtige Komponente in der Nährstoffversorgung von Plankton spielt und daher auch die Primärproduktion antreiben und somit auch zur Eutrophierung beitragen kann. In einer vierten Studie wurde das konservative gegen das nicht-konservative Verhalten von gelöstem organischem Material (DOM), welches DON und DOC (gelösten organischen Kohlenstoff) einschließt, entlang desselben Salzgehaltsgradienten getestet. Zusätzlich wurde der terrestrische Anteil vom DOM am Gesamt DOM mit Hilfe von  $\delta^{13}\text{C}$  Werten über Mischungsmodelle berechnet. Hier sprechen die Ergebnisse dafür das bis zu 83% des DOM in der Ostsee aus terrestrischen Quellen stammen und das der größte Anteil (>50%) vom DOM, das aus Flüssen stammt, in den Ästuaren und angrenzenden Küstengebieten abgebaut wird. Eutrophierung ist nicht nur ein Problem in der Ostsee, sondern weltweit und daher lassen sich einige Ergebnisse dieser Doktorarbeit auch auf andere Küstengebiete in den gemäßigten Breiten mit hohen Stickstofffrachten übertragen. Insgesamt ist es nötig die Rolle von Sedimenten in Bezug auf die Stickstoffentfernung zu überdenken und gelöste organische Substanzen als potentielle Nährstoffquellen zu berücksichtigen.

## 1. Introduction

### 1.1 Ecological problems of coastal eutrophication

At present, more than 40% of the world's ocean are strongly influenced by human activities (Halpern et al. 2008). Human population growth and its associated activities like fossil-fuel combustion, production of nitrogen fertilizers (Haber Bosch process) and cultivation of nitrogen-fixing legumes have increased the flux of nitrogen and phosphorus to aquatic and terrestrial ecosystems altering global cycles of both nutrients (Gruber and Galloway 2008, Galloway et al. 2003, 2004, Howarth 2006, 2008, Vitousek et al. 1997, Pickney et al. 2001). As a result of increasing nitrogen and phosphorus inputs the accumulation of organic matter by primary production has increased, which is defined as eutrophication (Nixon 1995). Selman et al. (2008) identified more than 400 areas worldwide which experienced symptoms of eutrophication (Fig. 1.1). Especially in coastal waters eutrophication has become the biggest pollution problem (e.g Vitousek et al. 1997, Syvitski et al. 2005, Rabalais 2002, Howarth et al. 2000), since rivers deliver large amounts of excess nutrients (Boyer et al. 2006, Dumont et al. 2005). It has been shown that nutrient river runoffs directly reflect human population density and activity in the watersheds (Peierls et al. 1991).




Figure 1.1: Map of 415 eutrophic and hypoxic coastal systems worldwide. The map was compiled by R. Diaz, M. Selman and Z. Sugg (<http://www.wri.org/map/world-hypoxic-and-eutrophic-coastal-areas>).

Besides increasing primary production, effects of eutrophication are enhancement of anoxia and hypoxia in deep and coastal waters, harmful algae blooms (HABs), decreasing water clarity and increased fluxes of  $N_2O$  to the atmosphere (Nixon 1995, Conley et al. 2009a, 2011, Galloway et al. 2003, Diaz and Rosenberg 2008). Even though hypoxia occurs naturally in many marine environments



(like fjords and deep basins (Gustafsson and Nordberg 2000, Zillen et al. 2008)), the so called dead zones have spread exponentially since the 1960s, approximately doubling the number each decade and affecting today more than 240.000 km<sup>2</sup> of the marine environment (Diaz and Rosenberg 2008). Also HAB's occur as a natural phenomenon, but over the last decades their extent has increased due to nutrient increases and/or shift in nutrient ratios (Anderson et al. 2002, Glibert et al. 2005, Heisler et al. 2008). These negative effects of eutrophication in turn can lead to loss or degradation of habitats with consequences to marine biodiversity and changes in ecosystem structure and function, such as cycling of elements and processing of pollutants (Vitousek et al. 1997, Rabalias 2002). Much of this eutrophication is driven by nitrogen (Howarth 2008). Due to the growing global demand in reactive nitrogen the pressure on coastal ecosystem may even rise in the future (ENA-book). To improve management and focus on strategies to resolve eutrophication a better understanding of coastal system dynamics is highly needed. This thesis will contribute in numerous ways, since it focuses on two major problems of eutrophication: dissolved organic matter (DOM) and N-retention and turnover in coastal waters.

## 1.2 Global nitrogen cycle

The largest pool of nitrogen is found in the atmosphere as dinitrogen gas (N<sub>2</sub>) that naturally only becomes biologically available through the process of N<sub>2</sub> fixation, the conversion of N<sub>2</sub> into organic nitrogen. The ability to fix N<sub>2</sub> is widespread in the marine environment, with cyanobacteria fixing 140 Tg N yr<sup>-1</sup> on a global scale (Brandes 2007, Gruber and Galloway 2008, Fig. 1.2). In terrestrial systems, N<sub>2</sub> fixation accounts for 100 Tg N yr<sup>-1</sup> (Fig. 1.2). With the onset of industrialization it became evident that the amount of reactive nitrogen (fixed nitrogen, Nr: including nitrate (NO<sub>3</sub><sup>-</sup>), nitrite (NO<sub>2</sub><sup>-</sup>), ammonium (NH<sub>4</sub><sup>+</sup>), and dissolved organic nitrogen (DON)) needed to be increased to sustain the growing population. With the invention of the Haber Bosch process, where nonreactive N<sub>2</sub> is converted to reactive NH<sub>3</sub> (ammonia) it was possible to meet the growing food demand. It has been estimated that nitrogen fertilizers are responsible for feeding nearly 50% of the world's population (Erisman et al. 2008). However, the nitrogen use efficiency for typical cultivated plants like rice, wheat and maize is typically below 40%, meaning that most of the N applied fertilizers is lost to the atmosphere via denitrification or to the aquatic environment via leaching. In the latter it is assimilated into biomass, and therefore fueling eutrophication (Canfield et al. 2010, Erisman et al. 2008). This movement of nitrogen from one effect to another as it cycles through environmental reservoirs is referred as the nitrogen cascade (Galloway et al. 2003). Besides the close links in the nitrogen cycle between watersheds, airsheds and the marine system the nitrogen cycle is closely linked to the carbon and phosphorus cycles (Fig. 1.2), which becomes obvious by the Redfield ratio, the molar stoichiometric relationship between C, N and P (106:16:1) in marine organic matter. Of

particulate relevance is the carbon cycle, since atmospheric  $\text{CO}_2$  has a central role in controlling climate (Sarmiento and Gruber 2002). The full scale of impacts of additional reactive nitrogen input globally still remains unknown but it is obvious that further intensification in agriculture, increasing energy use and population growth will continue to alter the terrestrial and marine nitrogen cycle and will amplify eutrophication in the future. Together fossil-fuel combustion, production of nitrogen fertilizers and increased cultivation of nitrogen-fixing legumes have doubled the inputs of reactive nitrogen compared to preindustrial times, now exceeding natural nitrogen sources, accounting to  $160 \text{ Tg N yr}^{-1}$ , (Galloway et al. 2003, Gruber and Galloway 2008, Fig. 1.2). Therefore understanding how human induced ecological change interacts with and affects the structure and functioning of large estuarine ecosystems adjoining coastal waters remains important (Paerl et al. 2006).




Figure 1.2: Natural and anthropogenic N fluxes on land and in the ocean. The tight coupling between the nitrogen cycles on land and in the ocean with those of carbon and phosphorus are shown. Blue fluxes denote natural (unperturbed) fluxes and red fluxes denote anthropogenic perturbation. Redrawn from Gruber and Galloway (2008)

Reactive nitrogen can be assimilated by most organisms and most of the organic nitrogen in the ocean is returned back to nitrate via remineralization to ammonium (ammonification), which is rapidly nitrified and therefore rarely occurs in significant concentrations in oxygenated habitats. The first step in nitrification is the oxidation of ammonium to nitrite by ammonia-oxidizing bacteria and archaea, the second step is the conversion of nitrite to nitrate by nitrite-oxidizing bacteria (Fig. 1.3). Nitrification produces the substrate for denitrification and anaerobic ammonium oxidation (anammox), the only two processes that remove nitrogen permanently from the system by producing  $\text{N}_2$ . Denitrification describes the conversion of nitrate into  $\text{N}_2$  gas through a series of

intermediates ( $\text{NO}_2^-$ ,  $\text{NO}$ , and  $\text{N}_2\text{O}$ ) (Knowles, 1982, Fig. 1.3). It takes place under suboxic and anoxic conditions when organic carbon (as electron donor) and nitrate (as electron acceptor) are both available (Knowles 1982, Seitzinger 1988). During anammox  $\text{NH}_4^+$  is anaerobically oxidized to  $\text{N}_2$  in the absence of organic matter (Dalsgaard et al. 2003, Kuypers et al. 2003, Fig. 1.3) and a recent examination revealed that it could possibly account for 30% to 50% of water column  $\text{N}_2$  production (Devol 2003). Apart from the reduction to  $\text{N}_2$  through these two processes, nitrate can also be reduced to ammonium during DNRA (Dissimilatory nitrate reduction to ammonium). The importance of DNRA in the water column nitrogen cycling is still widely unknown, but in the Oman Shelf DNRA was found to be an important mineralization pathway for organic matter (Jensen et al. 2011). Current global nitrogen budgets have shown that benthic denitrification and anammox can account for about 70% of the fixed nitrogen loss (Codispoti 2007). But if the nitrogen budget is in balance is still highly controversial (Gruber and Galloway 2008, Zehr and Kudela 2011).




Figure 1.3: The marine nitrogen cycle. The various chemical forms of nitrogen are plotted versus their oxidation state, where nitrate is the most oxidized N species, while ammonium and organic nitrogen comprise the most reduced species involved in the cycle. Processes shown in light grey occur in anoxic environments only (modified from Gruber 2008).

### 1.3 Dissolved organic matter as part of the nitrogen cycle

Dissolved organic matter (DOM) is a complex pool of organic molecules that pass a filter of nominal pore size 0.2–1  $\mu\text{m}$  (Hedges 2002). It consists of high molecular weight (HMW, weight > 1 kDa) and low molecular weight (LMW, weight < 1 kDa) compounds. HMW DOM includes proteins, dissolved combined amino acids, nucleic acids (DNA, RNA), and humic substances. LMW DOM consists of urea, dissolved free amino acids and amino sugars (Bronk 2002, Bermann and Bronk 2003). However, most compounds still remain chemically uncharacterized (Bronk 2002). Recent progress in analytical chemistry like the fourier transform ion cyclotron mass spectrometry (FT-ICR-MS) enable a characterization of dissolved organic matter at the molecular level in unprecedented detail (Koch et al. 2008, Dittmar and Paeng 2009), but still there is a gap in knowledge due to the chemical complexity of the compounds. According to Benner (2002) only 4-14% of DOM have been chemically characterized.

DOM in the marine environment can be from both allochthonous and autochthonous sources. Allochthonous sources include terrestrial runoff, leaching from plant detritus and soils into streams and rivers (Valiela et al. 1990, Tobias et al. 2001), sediments (Burdige 2002), groundwater (Santos et al. 2008) and atmospheric deposition (Duce et al. 2008, Cornell et al. 1995). Most of the allochthonous DOM has passed through several systems and chemical processes like phototransformation by sunlight, bacterial degradation in surface waters and soils as well as chemical changes due to increasing salinities before it reaches coastal regions. Autochthonous sources include release by primary producers (Bronk and Ward 1999, Stedmon et al. 2006) and bacteria (Ogawa 2001), excretion from micro and mesozooplankton (Steinberg et al. 2000), viral lysis of bacteria (Fuhrman 1999) and eukaryotic cells (Suttle 1994), and particle solubilization (Smith 1992) (Fig. 1.4). Main sinks of DOM are heterotrophic uptake, autotrophic uptake and abiotic photochemical decomposition (Bronk 2002) (Fig. 1.4). Small organic compounds like amino acids are taken up through permeases (membrane transport proteins), whereas larger compounds have to be broken down by extracellular hydrolytic enzymes (Anita et al. 1991, Bronk 2002, Mulholland and Lomas 2008 and references therein). For coastal areas terrestrial input via rivers dominates and highest concentrations of DOM are found close to the mouth of rivers (Stepanaukas et al. 2002, Feistel et al. 2008, Bronk 2002). In oceanic systems the major DOM sources are primary production and atmospheric input (Carlson 2002).

DOM is generally characterized in terms of carbon (DOC), nitrogen (DON) and/or phosphorus (DOP). DOC, DON and DOP can function largely independently within many ecosystems and are therefore discussed individually. In this thesis I will mainly focus on the DON components but also DOC will be discussed in face of its distribution and terrestrial background (see chapter 4 and 5 for details). DON

represents the largest pool of fixed nitrogen in most aquatic systems with concentrations decreasing from rivers to the open ocean (Bronk 2002). In the past DON has long been ignored as a potential N source, because it was thought to be mainly refractory. However, recent studies have shown that DON can also serve as an important N source for both phytoplankton and bacteria (Veuger et al. 2004, Seitzinger and Sander 1999, Middelburg and Nieuwenhuize 2000, Berg et al. 2001, Bronk et al. 2007), and thus contribute significantly to marine eutrophication (Berman 1997, Seitzinger and Sander 1997). For example, bioassay experiments in nine US rivers indicated that up to 23% of DON is bioavailable and that 43% of the consumed total dissolved nitrogen (TDN) by bacteria is taken up as DON (Wiegner et al. 2006). In general, the bioavailability of terrestrially derived DON is variable at 2-70% (Seitzinger and Sanders 1997, Stepanauskas et al. 2002, Veuger et al. 2004, Wiegner et al. 2006). It is speculated that the bioavailability and the composition of DON may depend on the source (McCallister et al. 2006). DON from anthropogenic sources seems to be more bioavailable than DON exported from forested regions and wetlands (Seitzinger et al. 2002). Especially in summer DON may significantly enhance primary production and its impact on the microbial community and contribution to eutrophication seems to be greater (Stepanauskas et al. 1999, 2002, Berg et al. 2003). Different compounds of DON, like urea or amino acids have been studied in detail (Bronk et al. 1998, Berman and Bronk 2003 and references therein), but studies on the bulk pool are rare (for details see chapter 4).

Like DON, DOC was also assumed to be mainly refractory and therefore unimportant as an energy source for microbes. However, studies in the last decades have shown that DOC is also highly bioavailable. Amon et al. (2001) demonstrated that 30% of fresh, algal-derived DOM was bioavailable in form of DOC for bacteria. Terrigenous dissolved organic carbon was shown to be relatively rapidly remineralized on the continental shelf of the Arctic Ocean (Letscher et al. 2011). In the world ocean approximately 97% of all organic carbon exists in the dissolved phase, which is comparable to the amount of carbon dioxide in the atmosphere (e.g. Siegenthaler and Sarmiento 1993). Every year around 0.25 Gt C are transported as DOC via the rivers to the coastal oceans and a relatively high proportion is degraded after mixing with seawater with turnover rates ranging from days to years (Cauwet et al. 2002). Additionally, only a small fraction of the organic matter within the ocean appears to be land derived (Hedges et al 1997, see chapter 5 for more details). Even though DOC and DON cycling is partly intertwined, it has been shown that DON can be preferentially recycled to avoid nitrogen limitation (Thomas et al. 1999). But still the dynamics of specific compounds of DOM and much less of the bulk pool remains unclear and therefore tracing DOM with stable isotopes can help understanding its cycling (see chapter 4 and 5 for more details).




Figure 1.4: Conceptual diagram of autochthonous sources and sinks of DON in the marine environment. DPA stands for dissolved primary amines (Bemann and Bronk 2003, Bronk 2002).

#### 1.4 Stable isotopes as a tool to unravel the nitrogen cycle

To assess nitrogen turnover processes and sources, nitrogen and oxygen isotopes in natural abundance and nitrogen isotopes in artificial labeling experiments were used in many aquatic studies (e. g. Dugdale and Wilkerson 1986, Liu & Kaplan 1989, Brandes et al. 1998, Sigman et al. 1999, Middelburg and Nieuwenhuize 2000, Voss et al. 2000, Sutka et al. 2004, Deutsch et al. 2006, Casciotti et al. 2007, Dähnke et al. 2008, Granger et al. 2008, see chapter 3 and 4 for details). Isotopes are atoms with the same number of protons but different number of neutrons. Stable isotopes are defined as those that are energetically stable and do not decay. Nitrogen has two stable isotopes  $^{14}\text{N}$  and  $^{15}\text{N}$ , while oxygen has three,  $^{16}\text{O}$ ,  $^{17}\text{O}$  and  $^{18}\text{O}$ . In this study  $^{17}\text{O}$  is neglected due to its very small abundance in natural sources, but for the mass-dependent  $^{16}\text{O}/^{17}\text{O}/^{18}\text{O}$  relationship a correction was always made for  $^{17}\text{O}$ . Isotope ratios are expressed in the delta notation ( $\delta^{15}\text{N}$  relative to atmospheric  $\text{N}_2$  and  $\delta^{18}\text{O}$  relative to Vienna Standard Mean Ocean Water (VSOW)):

$$\delta_{\text{sample}} (\text{‰}) = (R_{\text{sample}}/R_{\text{reference}}) \times 1000$$

where R is  $^{15}\text{N}/^{14}\text{N}$  or  $^{18}\text{O}/^{16}\text{O}$  ratio of sample and reference, respectively. Stable isotopes values are altered by both equilibrium processes (like evaporation of water) or by kinetic fractionation processes which tend to partition light isotopes ( $^{14}\text{N}$  and  $^{16}\text{O}$ ) from heavier ones ( $^{15}\text{N}$  and  $^{18}\text{O}$ ) (Kendall 1998). As a consequence, throughout the course of a biochemical reaction, the substrate being consumed becomes progressively enriched with the heavier isotope, while the resultant product becomes relatively lighter. The extent to which a biological transformation fractionates between light and heavier isotopes is given by the isotopic effect,  $\epsilon$  (Tab. 1.1).

This value is calculated from the integrated expression of the progress of the reaction according to the Rayleigh model:

$$\delta_{\text{reactant}} = \delta_{\text{initial}} - \varepsilon[\ln(f)]$$

where  $f$  is the fraction of reactant remaining (nitrate/nitrate<sub>initial</sub>),  $\delta_{\text{initial}}$  is the  $^{15}\text{N}/^{14}\text{N}$  or  $^{18}\text{O}/^{16}\text{O}$  ratio of initial reactant pool, and  $\varepsilon$  is the kinetic isotope effect of the transformation. In practice,  $\varepsilon$  is the negative slope of the linear relation of  $\delta^{15}\text{N}$  or  $\delta^{18}\text{O}$  vs. the natural logarithm of the fraction of the reactant remaining.

Table 1.1: Fractionation factors ( $\varepsilon$ ) for N cycle processes.

Reaction	$^{15}\varepsilon$	$^{18}\varepsilon$	References
Nitrate assimilation ( $\text{NO}_3^- \rightarrow \text{PN}$ )	~5‰	$^{18}\varepsilon = ^{15}\varepsilon$	Wu et al. 1997, Sigman et al. 1999, Altabet 2001, Granger et al. 2004
Nitrification ( $\text{NH}_4^+ \rightarrow \text{NO}_2^-$ )	14-38‰	-	Casciotti et al. 2003
Water column denitrification ( $\text{NO}_3^- \rightarrow \text{N}_2$ )	22-30‰	$^{18}\varepsilon = ^{15}\varepsilon$	Brandes et al. 1998, Altabet et al. 1999, Voss et al. 2001
Sediment denitrification ( $\text{NO}_3^- \rightarrow \text{N}_2$ )	<3‰	$^{18}\varepsilon = ^{15}\varepsilon$	Brandes and Devol 1997, Lehmann et al. 2004
$\text{N}_2$ fixation ( $\text{N}_2 \rightarrow \text{NH}_4^+$ )	-1,5-0‰	~0‰	Capone et al. 1997, Minagawa and Wada 1986
Ammonification ( $\text{N}_{\text{org.}} \rightarrow \text{NH}_4^+$ )	~0‰	-	Kendall 1998

#### 1.4.1 Tracking nitrogen in coastal areas

New insights into the nitrogen cycle have been gained from studies that have used stable isotopes in artificial labeling experiments and natural abundances. For example studies using  $^{15}\text{N}/^{14}\text{N}$  isotopic tracers examined the linkages in the cycle of new nitrogen input into the euphotic zone of the ocean, its utilization by phytoplankton, and transport to the deep sea (Altabet and Deuser 1985, Altabet and McCarthy 1985, 1986, Altabet 1988, 1989, Voss et al. 1996). Further on, the distribution of nitrogen

isotopes within marine ecosystems can provide a record of the sources of nitrogen supporting biological production and the major pathways and mechanisms moving nitrogen through the biota (Montoya 2008).

Because about 88% of the oceanic fixed nitrogen pool consists of nitrate (Gruber 2008) many studies over the last decades have used  $^{15}\text{N}$  isotopes in nitrate to investigate sources and transformation processes (Brandes and Devol 2002 and references therein). Even though, measurements of  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  in nitrate in fresh water samples were possible (Silva et al. 2000) the invention of the denitrifier method (Sigman et al. 2001, Casciotti et al. 2002, see chapter 3 for details) enabled researchers to investigate the influence of both cycling and mixing of multiple sources in marine water samples with low nitrate concentrations (e.g. Wankel et al. 2006, Dähnke et al. 2010, Sebilo et al. 2006, Sigman et al. 2003). The advantage of the additional measurement of the O isotopes in nitrate is that now processes overprinting each other when looking at nitrogen isotopes in nitrate can be distinguished. For example, Sigman et al. (2003) were able to distinguish between water column and sedimentary denitrification in the Santa Barbara Basin by using the dual stable isotopes of nitrate. They could furthermore show that sedimentary denitrification accounts for more than 75% of the nitrate loss within that area.

The most significant processes in rivers and coastal areas that cause isotopic fractionation are assimilation, nitrification, and denitrification. Assimilation as well as denitrification results in an increase of  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  values in nitrate (Kendall 1998) because plankton and microbes preferentially consume isotopically light nitrate ( $^{14}\text{N-NO}_3^-$  and  $^{16}\text{O-NO}_3^-$ ) (Mariotti et al. 1988, Lui & Kaplan 1989, Kendall 1998, Voss et al. 2001, Lehmann et al. 2003). The  $\delta^{15}\text{N}/\delta^{18}\text{O}$  ratio of the remaining nitrate is supposed to be close to 1:1 during assimilation (Granger et al. 2004). The same holds true for denitrification in marine environments (Sigman et al. 2005, Granger et al. 2004), whereas in freshwater ecosystems the ratio seems to be 1:0.5 (Böttcher et al. 1990, Lehmann et al. 2003). But also anomalies from the 1:1  $\delta^{15}\text{N}/\delta^{18}\text{O}$  relationship can be found in the marine environment (Sigman et al. 2005, see chapter 3 for details). During nitrification the development of  $\delta^{15}\text{N-NO}_3^-$  and  $\delta^{18}\text{O-NO}_3^-$  is decoupled. Nitrification adds isotopically depleted nitrate to the nitrogen pool because nitrifiers preferentially take up isotopically light ammonium (Kendall 1998). In addition to the N isotope effects, also O isotope effects are involved with the nitrification process (Buchwald and Casciotti 2010, Casciotti et al. 2010). New O atoms are added from dissolved oxygen ( $\text{O}_2$ ) and water ( $\text{H}_2\text{O}$ ) pools, independently of other N cycle processes. While  $\text{O}_2$  is incorporated during the oxidation of  $\text{NH}_3$  to  $\text{NH}_2\text{OH}$ ,  $\text{H}_2\text{O}$  is incorporated during the oxidation of  $\text{NH}_2\text{OH}$  to  $\text{NO}_2$  and  $\text{NO}_3^-$  (Buchwald and Casciotti 2010, Casciotti et al. 2010). It was suggested that the  $\delta^{18}\text{O-NO}_3^-$  signature from nitrification is dominated by the water  $\delta^{18}\text{O}$  signal, because 5 out of 6 oxygen atoms in nitrate originate from the water (Casciotti et al. 2002, Sigman et al. 2005). But this only holds true when



exchange and fractionation of oxygen isotopes during nitrification are minimal (Casciotti et al. 2011). Casciotti et al. (2010) and Buchwald and Casciotti (2010) could show that in addition to variations in the oxygen isotope value of the O atom donors ( $O_2$  and  $H_2O$ ), the oxygen isotope value of newly produced nitrate is affected by O isotopic exchange and fractionation. This makes it even more complicated to define a fractionation factor for nitrification and additional work is needed to fully characterize the O isotopic systematics for nitrification (Casciotti et al. 2011).




Figure 1.5: Ranges of isotopic compositions of nitrate ( $\delta^{15}N$  and  $\delta^{18}O$ ) for major nitrate sources and the expected trend for the isotopic composition undergoing denitrification (adopted from Mayer et al. 2002, Kendall et al. 2007)

The isotope values ( $\delta^{18}O$ - and  $\delta^{15}N$ - $NO_3^-$ ) resulting from various turnover processes can be used to attribute nitrate to specific sources (Fig. 1.5, Boyer et al. 2002, Mayer et al. 2002, Voss et al. 2006, Deutsch et al. 2006). Enriched  $\delta^{15}N$ - $NO_3^-$  values indicate sewage input and/or input of fertilizers in agriculture runoff ranging from 7 and 25‰ (summarized in Kendall et al. 2007). Compared to nitrate from natural sources anthropogenic nitrate is enriched in  $^{15}N$  by various processes like ammonia volatilization and denitrification in soils or aquifers. The stable nitrogen isotopic composition of atmospherically deposited nitrogen is highly variable and can be influenced for example by fossil fuel composition or agricultural activities (Kendall 1998, Mayer et al. 2001, Mayer et al. 2002). Synthetic fertilizers and nitrate in natural soils are isotopically distinct from N in the atmosphere and  $\delta^{15}N$ - $NO_3^-$  and  $\delta^{18}O$ - $NO_3^-$  vary between 0 to 3 and  $22 \pm 3$ ‰ and -3 to 5 and 0 to 14‰, respectively (Kendall 1998). Therefore the isotopic composition of nitrate is a powerful tool to determine both nitrogen transformation processes and its origin (see chapter 3 for details).

### 1.5 The Baltic Sea and its sensitivity towards eutrophication

The Baltic Sea is one of the most eutrophied coastal seas in the world (Cloern 2001) and anthropogenic eutrophication effects are well studied (Elmgren 2001). Every year about 1000 kt of nitrogen are entering the Baltic Sea from various sources like rivers, N<sub>2</sub> fixation, atmospheric deposition and point sources (Tab. 1.2). In chapter 2 detailed background informations on the Baltic Sea are given. Therefore, only a brief summary in this section about the Baltic Sea is given to highlight why it is an excellent site for studying nitrogen transformation processes in face of eutrophication.

Table 1.2: Nitrogen inputs to the Baltic Sea

Input pathway	Baltic Sea N input [kt N yr <sup>-1</sup> ] in 2000	Reference
Riverine	745	Helcom 2005
Atmospheric	264	Helcom 2005
N <sub>2</sub> fixation	180 to 430	Rolff et al. 2008, Wasmund et al. 2005

The Baltic Sea is a shallow, intra-continental, brackish sea and covers an area of 377.000 km<sup>2</sup> with a coastline of 8000 km (Sweitzer et al. 1996). Its topography is characterized by a series of basins (e.g. Arkona, Bornholm and Gotland Basins). A positive water balance creates a strong salinity gradient from 2 in the northernmost Gulf of Bothnia to over 20 in the Kattegat and Danish Straits and a permanent halocline separates the deep, saltier water from the surface water in the western and central Baltic Sea. The bottom water of the Baltic proper is only replaced by intermittent inflows of denser, oxygenated water through the Danish Straits from the North Sea. Additionally due to the limited water exchange with the North Sea and long residence times of approximately 30 years, stagnation periods of up to 16 years enhance the natural occurrence of anoxic bottom waters (Elmgren 2001, Conley et al. 2002). Anthropogenic driven eutrophication has increased both the spatial extent and intensity of hypoxia in the deep basins of the Baltic Sea (Conley et al. 2009, Zillen et al. 2008). Recently a study could show that over 115 coastal sites in the Baltic Sea experienced hypoxia, which implies that over 20% of all worldwide known hypoxic sites are in the coastal zone of the Baltic Sea (Conley et al. 2011). The consequences of hypoxia on the biota are mentioned in section 1.1 but also the consequences on the nutrient biogeochemical cycles of the Baltic Sea are substantial leading to increased phosphorus release from the sediments (Mort et al. 2010) and internal loading and circulation of nutrients are intensified. Additionally, eutrophication has led to an

increase in the abundance of cyanobacteria and other phytoplankton blooms in the Baltic Sea (Finni et al. 2001, HELCOM 2005, Vahtera et al. 2007).

Another characteristic of the Baltic Sea is that the drainage basin of the Baltic Sea is four times larger than the sea itself and is populated by 85 million people (Sweitzer et al. 1996). The population is heterogeneously distributed with a gradient from high population density in the south ( $>500$  inhabitants/km<sup>2</sup>) to low population density in the northern part ( $< 10$  inhabitants/km<sup>2</sup>) (Lääne et al. 2005). In the north the land cover is dominated by boreal forests whereas the southern catchment is heavily cultivated. This results in higher riverine inputs of nutrients from the southern part of the catchment (HELCOM 2004, HELCOM 2009, Humborg et al. 2007, Voss et al. 2011). For the Baltic Sea rivers play a crucial role in the total input of nitrogen, accounting for approximately 75% of the total nitrogen input to the Baltic Sea (HELCOM 2005, Rolff et al. 2008, Table 2). The rivers Neva, Vistula, Daugava, Nemunas and Oder supply over 50% of the total nitrogen influx mainly in form of dissolved inorganic nitrogen (DIN) (HELCOM 2004). It is well known that concentrations decrease rapidly offshore in the Baltic Sea but the nitrogen removal processes which lead to the loss of nitrate are still not fully characterized in the estuaries and adjacent coastal areas (see chapter 2 and 3 for details).

To solve the ecological problems of the Baltic Sea, in 1974 seven Baltic coastal states formed the Helsinki Commission (HELCOM). In 2007 the Helsinki Commission (now nine Baltic coastal states) adopted the Baltic Sea Action Plan (BSAP). In the BSAP clear nutrient reduction goals have been allocated to the riparian countries and the main goal is that the Baltic Sea is supposed to be unaffected by eutrophication by 2021. The same is claimed in the European Water Framework Directive (WFD 2000), which implements that in surface waters like coastal water bodies, including the Baltic Sea, good ecological and chemical conditions are to be reestablished. Although recent data indicate a decrease in nitrogen runoff (Wulff et al. 2009), eutrophication remains the single greatest threat to the Baltic Sea environment (HELCOM 2009, Andersen et al. 2011). In a recent study the HELCOM still claims 176 areas out of 189 to be affected by eutrophication (Andersen et al. 2011).

It has been shown that denitrification as the main process that removes nitrogen permanently from the system is estimated to account for 48-73% of external nitrogen removal (Deutsch et al. 2010). Additionally, Hietanan et al. 2008 could show that anammox can contribute 10 to 15% to the total N<sub>2</sub> production in the Baltic Sea. But still the nitrogen budget in the Baltic Sea is not balanced. Additionally, two recent studies suggest that the removal capacity may even decrease when DNRA becomes the most important pathway under hypoxia (Jäntti et al. 2011, Jäntti et al. in press). Therefore studying nitrogen turnover and removal processes in the Baltic Sea is of great importance to make sure that the nitrogen removal capacities are not lost, besides the effort to reduce nitrogen loads.

### 1.6 Aim of the study

This thesis is part of the international BONUS (Baltic Organization Network for Funding Science) project AMBER (Assessment and Modelling of Baltic Ecosystem Response), which focuses on the implementation and application of an Ecosystem Approach to Management (EAM) to the Baltic Sea in face of eutrophication and climate change focusing on the coastal ecosystem. The aim of the present study is to characterize nitrogen turnover processes in the Baltic Sea and to contribute to the existing knowledge on the role of nitrogen in coastal areas. The study is based on the one hand on in situ rate measurements in artificial labeling experiments and on the other hand on the analyses of natural stable nitrogen, oxygen and/or carbon isotopes in different pelagic compounds like nitrate or particulate organic matter (POM).

In chapter 2 the existing knowledge on the historical progression of eutrophication based on long term data sets is summarized. Additionally, the sensitivity of the Baltic Sea towards eutrophication and future scenarios from climate related change in the Baltic Sea region are examined. Regional differences that lead to major open questions concerning the nitrogen cycle in the Baltic Sea are elucidated. The goals of chapter 3 are to trace the isotopic signature of riverine nitrate in the coastal zone of the Baltic Sea and to determine main nitrate removal processes. Thus nitrate from different sources, including the Oder River in the south, the Nemunas River in the north east and the central Baltic Sea are analyzed by means of dual isotope analysis. Additionally,  $^{15}\text{N}$  nitrate tracer techniques were used to quantify the nitrate uptake by in situ measurements. Also the influence of permeable sediments on the N-cycling in the overlying bottom water is studied. This part is based on data from a spring cruise in 2009, when nitrate loads and river runoff reach their annual high. In chapter 4 and 5 the fate of organic matter (DON and DOC) is characterized by means of  $^{15}\text{N}$  and  $^{13}\text{C}$  tracer techniques along a salinity gradient ranging from 34 to 2. Turnover rate calculations and end-member-mixing-models were used to show the high dynamics of the bulk DOM pool. These studies were performed during summer to account the fact that DON can be a nutrient source, when inorganic nitrogen is limited. Chapters 2 to 5 are presented in a manuscript-like structure and a detailed statement on my contributions to the manuscripts are given at the end of the thesis. In chapter 6 the results of this thesis are summarized and a short outlook is presented.

## 2. History and scenarios of future development of Baltic Sea eutrophication

M. Voss, J. Dippner, C. Humborg, F. Korth, T. Neumann, J. Hürdler, G. Schernewski, M. Venohr

Estuarine, Coastal and Shelf Science, 92(3), pp. 307-322

### 2.1a Abstract

Nutrient loads from watersheds, atmospheric deposition, and cyanobacterial nitrogen fixation have led to eutrophication in the Baltic Sea. Here we give the historical evolution of this, detail some of the specific eutrophication features of the Baltic Sea, and examine future scenarios from climate related changes in the Baltic Sea region. We distinguish northern and southern regions of the Baltic Sea. The northern watersheds have sub-polar climate, are covered by boreal forest and wetlands, are sparsely populated, and the rivers drain into the Gulf of Bothnia. The southern watersheds have a marine influenced temperate climate, are more densely populated and are industrially highly developed. The southern areas are drained by several large rivers, including the representative Oder River. We compare these regions to better understand the present and future changes in Baltic Sea eutrophication.

Comparing the future projections for the two regions, we suggest that in addition to changes in nutrient inputs, increased temperature and precipitation are likely to become important forcings. Rising temperature may increase release of dissolved organic matter (DOM) from soils and may alter the vegetation cover which may in turn lead to changed nutrient and organic matter input to the Baltic Sea. For the southern Oder River catchment a model study of nutrient input is evaluated, MONERIS (Modelling Nutrient Emissions in River Systems). The strong correlation between precipitation, flow and nutrient discharge indicates a likely increase in nutrient concentrations from diffuse sources in future. The nutrients from the Oder River are modified in a lagoon, where removal processes change the stoichiometry, but have only minor effects on the productivity. We suggest that the lagoon and other nearshore areas fulfil important ecological services, especially the removal of large quantities of riverine nitrogen but at the same time are threatened systems due to increasing coastal hypoxia.

### 3. Nitrogen turnover during spring outflow from the nitrate-rich Curonian and Szczecin lagoon using dual isotopes in nitrate

F. Korth, I. Liskow, B. Fry and M. Voss

Submitted to Marine Chemistry

#### 3.1 Abstract

Coastal zones which receive high nitrogen loads from rivers often suffer substantially from eutrophication. This is true for the Baltic Sea, which is well-known for its critical eutrophication status especially along the coastlines. Although the nutrient concentrations decrease rapidly offshore, the uptake and turnover processes near-shore are not well understood. The Rivers Oder and Nemunas, the second and third largest nitrogen contributors, drain into the Szczecin and Curonian lagoon, respectively, before they enter the coastal area of the Baltic Sea. During peak outflow, in March 2009 nutrient concentrations, nitrate uptake rates and dual isotopes ( $\delta^{15}\text{N-NO}_3^-$  and  $\delta^{18}\text{O-NO}_3^-$ ) in nitrate were measured in the outflows of the lagoons to characterize nitrate turnover processes and its fate in the coastal zone. In the Curonian lagoon outflow the isotopic signature is dominated by mixing, whereas in the Szczecin lagoon outflow the isotope values are influenced by the ongoing spring phytoplankton bloom. Nitrate assimilation is indicated in the surface waters of the Szczecin lagoon outflow by a parallel enrichment of  $^{15}\text{N}$  and  $^{18}\text{O}$ . In the near bottom waters denitrification seems to be the prevalent process which generates the isotopic signal of nitrate. There a deviation from the 1:1  $\delta^{18}\text{O-NO}_3^-$  to  $\delta^{15}\text{N-NO}_3^-$  relationship usually associated with denitrification was found. This 1.3:1 ratio suggests that denitrification is not only fueled by nitrate fluxes from the water column into the sediments, but also from nitrate derived from remineralization of particulate matter and coupled nitrification/denitrification. Moreover, the fractionation factors of  $^{15}\epsilon$  of 9.9‰ and  $^{18}\epsilon$  of 10.1‰ in near bottom waters infer that the isotopic enrichment from sedimentary denitrification may be higher in sandy sediment than in muddy sediments.

#### 4. Uptake of dissolved organic nitrogen by size-fractionated plankton along a salinity gradient from the North Sea to the Baltic Sea

F. Korth, B. Deutsch, I. Liskow, M. Voss

Biogeochemistry (DOI 10.1007/s10533-011-9656-1)

##### 4.1 Abstract

The Baltic Sea is known for its ecological problems due to eutrophication caused by high nutrient input via nitrogen fixation and rivers, which deliver up to 70% of nitrogen in the form of dissolved organic nitrogen (DON) compounds. We therefore measured organic nitrogen uptake rates using self produced  $^{15}\text{N}$  labeled allochthonous (derived from *Brassica napus* and *Phragmites sp.*) and autochthonous (derived from *Skeletonema costatum*) DON at twelve stations along a salinity gradient (34 to 2) from the North Sea to the Baltic Sea in August/September 2009. Both labeled DON sources were exploited by the size fractions 0.2-1.6  $\mu\text{m}$  (bacteria size fraction) and >1.6  $\mu\text{m}$  (phytoplankton size fraction). Higher DON uptake rates were measured in the Baltic Sea compared to the North Sea, with rates of up to 1213  $\text{nmol N L}^{-1} \text{h}^{-1}$ . The autochthonous DON was the dominant nitrogen form used by the phytoplankton size fraction, whereas the heterotrophic bacteria size fraction preferred the allochthonous DON. We detected a moderate shift from >1.6  $\mu\text{m}$  plankton dominated DON uptake in the North Sea and central Baltic Sea towards a 0.2-1.6  $\mu\text{m}$  dominated DON uptake in the Bothnian Bay and a weak positive relationship between DON concentrations and uptake. These findings indicate that DON is an important component of plankton nutrition and can fuel primary production. It may therefore also contribute substantially to eutrophication in the Baltic Sea especially when inorganic nitrogen sources are depleted.

## 5. Tracing inputs of terrestrial high molecular weight dissolved organic matter within the Baltic Sea Ecosystem

B. Deutsch, A. Alling., C. Humborg, F. Korth, C. M. Mörtz

Biogeosciences, 9, 4465-4475, 2012

doi:10.5194/bg-9-4465-2012

### 5.1 Abstract

To test the hypothesis whether high molecular weight dissolved organic matter (HMW-DOM) in a high latitude marginal sea is dominated by terrestrial derived matter, 10 stations were sampled along the salinity gradient of the central and northern Baltic Sea and were analyzed for concentrations of dissolved organic carbon as well as  $\delta^{13}\text{C}$  values of HMW-DOM. Different end-member-mixing models were applied to quantify the influence of terrestrial DOM and to test for conservative versus non-conservative behavior of the terrestrial DOM in the different Baltic Sea basins. The share of terrestrial DOM to the total HMW-DOM was calculated for each station, ranging from 43 to 83%. This shows the high influence of terrestrial DOM inputs for the Baltic Sea ecosystem. The data also suggest that terrestrial DOM reaching the open Baltic Sea is not subject to substantial removal anymore. However compared to riverine DOM concentrations, our results indicate that substantial amounts of HMW-DOM (> 50%) seem to be removed near the coastline during estuarine mixing. A budget approach yielded residence times for terrestrial DOM of 2.8, 3.0, and 4.5 yr for the Bothnian Bay, the Bothnian Sea and the Baltic Proper.



## 6. Conclusions and Perspectives

Currently over 40% of the world's population live within 100 kilometers of the coast and since the onset of industrialization the world population has grown to 7 billion people (The state of world population 2011) and will continue to increase. The UK Food and Agriculture Association estimates that the world population will increase to 8.9 billion, by 2050 meaning that global agriculture must increase even further in the next 30 years to sustain this type of population growth. As population density and economic activity in the coastal zone increases, pressures on coastal ecosystems will increase. Also changes in diet are of great importance. Eriksson Hägg et al. 2010 hypothesized that increased protein consumption in the eastern European countries could lead to 16 to 32% increased total nitrogen fluxes to the Baltic Sea making it even more difficult to reduce the nitrogen load by 135000 tons and phosphorous loads by 15250 tons as required by the Baltic Sea Action Plan (BSAP) for the year 2021. Also Krämer et al. (2011) estimated that agricultural nitrogen loads to the Oder Lagoon could increase by as much as 23% in the future considering aspects like cultivation of energy maize and increased animal stocks. Therefore, current and expected future human alterations to the nitrogen cycle make it necessary to understand how nitrogen is cycling through watersheds to assess the consequences of increasing anthropogenic nitrogen inputs.

In the framework of this thesis, the fate of dissolved organic and dissolved inorganic nitrogen were studied in detail in the Baltic Sea. The results indicate that not only inorganic but also organic nitrogen serves as an important nutrient source for phyto- and bacterioplankton and that it is mainly degraded in estuaries and near the coastline. In general, continental shelf ecosystems, such as the Baltic Sea, are characterized by strong gradients in nutrients and organic matter, which are mainly maintained by the continuous supply through river runoff. These gradients imply an intensive nutrient and organic matter cycling within the coastal areas, which contribute a major part to the annual nitrogen demand (e.g. Costanza et al. 1997, Middelburg and Nieuwenhuize 2001, Seitzinger et al. 2006). In the Baltic Sea pronounced regional differences are found according to the land use and population density. In the northern watersheds, which are covered by boreal forests and are sparsely populated, nitrogen discharges from the rivers are lower than in the southern region, where watersheds are densely populated and agriculture and industry are highly developed (Fig.6.1). Additionally, the total nitrogen (TN) composition varies (Stepanaukas et al. 2002). In the northern part organic nitrogen dominates, whereas in the southern part nitrate is the main component of the total nitrogen pool. Moreover, DON from anthropogenic sources is more bioavailable than DON exported from natural regions like forests or wetlands (Seitzinger et al. 2002). Congruently, results of this thesis indicate that the potential of plankton to utilize DON is higher in the southern part of the Baltic Sea, where anthropogenic loads are high and nitrogen limitation occurs (Graneli et al. 1990).

Pronounced differences were found between the North Sea and the Baltic Sea regarding the magnitude of DON uptake rates. In the Baltic Sea, where riverine DON loads are high, DON uptake rates were an order of magnitude higher than in the North Sea. This is the first study where bulk dissolved organic nitrogen (DON) uptake rates were determined by imitating a natural DON source with  $^{15}\text{N}$  labeled DON instead of using single DON compounds such as urea or amino acids. Stepanauskas et al. 2002 showed that around 30% of terrestrial derived DON was available to bacteria during bioassay experiments in the Baltic Sea. This thesis adds to the knowledge that besides bacteria also phytoplankton utilizes DON (either directly or following photodegradation or bacterial breakdown).




Figure 6.1: The Baltic Sea and its catchment area, which is colored according to the land use. Concentrations of total nitrogen (TN) ( $\mu\text{mol l}^{-1}$ ) are given for 35 rivers and the percentage of dissolved inorganic nitrogen (DIN), dissolved organic nitrogen (DON), and particulate nitrogen (PN) (sources Baltic Nest institute and Stepanauskas et al. 2002).

However, riverine loads are the most important allochthonous DON sources. From results of this thesis it was calculated that the share of terrestrial DOM to the total DOM pool in the Baltic Sea is high, ranging from 43 to 83%. Turnover rate calculations showed that terrestrial derived DON must be mainly degraded in near proximity of the coast and that only more refractory compounds remain. Additionally, in this thesis it was demonstrated that also DOC is an important energy source which is also mainly degraded in river estuaries and in the coastal area. The fate of carbon and nitrogen in organic matter can be coupled or decoupled (Veuger and Middelburg 2007, Engel et al. 2002, Bode et al. 2004). Especially under nitrogen limitation the utilization of DON seems to become important. Therefore future studies should trace both DON and DOC to investigate carbon-nitrogen couplings or decoupling. In order to quantify the loss of DON/DOC in the coastal zone, future studies should include measurements of bulk DON/DOC uptake rates along a DOM concentration gradient from the river to the open sea. Since the composition and reactivity of DON depend on its source and DON of different rivers is chemically distinct (McCallister et al. 2006) studies on the fate of DON should be performed in different river systems and over longer time periods, because the reactivity of DON can also vary over the season and even during peak outflow events (Stepanaukas et al. 2000, Wiegner et al. 2009, Bode et al. 2004).

In some regions already today dissolved inorganic nitrogen reductions can be observed due to improved fertilizer, livestock management and/or advanced wastewater treatment (Carstensen et al. 2006), but the productivity has not decreased, which is often attributed to internal nutrient loadings and recycling (Soetaert et al. 2006, Philippart et al. 2007). In the Oder River, one of the most eutrophied rivers of the Baltic Sea, a slight decrease in total nitrogen loads has been observed (Savschuk and Wulff 2009) due to inorganic nitrogen reductions, however around 32% of the nitrogen load is in form of DON and no reduction in DON can be seen. Since this thesis has shown that DON can be an important source for primary production it becomes even more necessary to focus as well on DON dynamics besides DIN and to have a closer look on DON sources and sinks. To evaluate its potential importance in ecosystem-wide nutrient budgets DON should be included as an extra nutrient sources in physical-biogeochemical models. Additionally, as suggested in chapter 2 climate change may lead to massive release of DOC and DON in boreal and arctic rivers since permafrost areas will start to thaw. If this additional DON and DOC will affect the productivity in coastal seas, is not clear yet (Hood and Scott 2008).

Further on, results from this thesis have shown that during cyanobacteria blooms DON release and uptake are closely coupled. Especially under the aspect that the frequency and magnitude of cyanobacteria blooms have increased in the Baltic Sea (Finni et al. 2000, Vahtera et al. 2007) and that climate change is a potential catalyst for the expansion of these blooms (Pearl and Huisman 2008, Meier et al. 2011) it becomes obvious that the understanding of DON dynamics is important to

understand how this pool will respond to anthropogenic perturbations and climate change. However, still numerous questions remain concerning dissolved organic substances and its role in the nitrogen cycle. Therefore further studies with  $^{15}\text{N}$  tracer techniques, for example by combining nitrogen concentration and stable isotope ratio measurements of DON with the “denitrifier method” are envisioned which can give more insights into processes (Schlarbaum et al. 2010, Knapp et al. 2005, 2011). With the combination of these methods it already have been shown in the North Atlantic and the North Pacific Ocean, as well as in coastal areas like the North Sea, that DON is actively participating in the nitrogen cycle of each region (Schlarbaum et al. 2010, Knapp et al. 2011).

Like organic substances also inorganic nutrients are known to decrease rapidly offshore specifically in the southern Baltic Sea, where the anthropogenic nitrogen isotopic signature is rapidly replaced by the one of nitrogen fixing organisms (Voss et al. 2000, 2005, 2006, HELCOM 2007). However the processes which influence the isotope values in the transition zone of riverine and marine water remained unclear. Therefore measurements of dual isotopes in nitrate were performed for the first time in the outflows of the Nemunas and Oder Rivers, two of the main nitrogen contributing rivers to the Baltic Sea, during peak outflow season, when nitrate loads are highest. Results indicate mixing in the surface water and nitrate assimilation, whereas in permeable sediments, as they are found in the Oder outflow, nitrate is denitrified directly or removed via coupled nitrification/denitrification. So far, permeable sediments were mainly thought to contribute little to the biogeochemical cycling due to its low carbon content thus biogeochemical research has focused on muddy shelf sediments. This study adds to the growing knowledge that permeable sediments can be important sites for nitrogen removal (Jahnke et al. 2005, Janssen et al. 2005). A recently published study by Gao et al. (2012) demonstrated for the North Sea that permeable, sandy sediments, which account for 58-70% of the continental shelf area (Emery 1968), are important nitrogen sinks. They estimated an annual nitrogen removal rate of  $745 \pm 109 \text{ mmol N m}^{-2} \text{ yr}^{-1}$  for these sediments. Future studies should consider the potential of permeable sediments to regulate the flow of nitrogen at the land–sea boundary in estuaries where nitrate and fresh organic matter loads are high. Consequently the contribution of permeable sediments to the global nitrogen loss budget need to be reevaluated.

So far denitrification rate measurements in permeable sediments in the Baltic Sea are rare (Deutsch et al. 2010), and have not been performed in the estuaries of the Oder and/or Nemunas River. Future studies should focus on the determination of denitrification rates in these permeable sediments to estimate the amount of nitrate lost in the estuaries and the adjacent coastal zone. Rate measurements should be performed seasonally to determine which major factors are controlling denitrification rates, since the interaction of parameters like substrate availability, temperature, and/or oxygen concentration is still under debate (Seitzinger 1988). Concerning denitrification rate

measurements the sample resolution in space and time is relatively low as collecting and incubation by using the isotope pairing technique is time consuming (Nielsen 1992, Risgaard-Petersen et al. 2003). Furthermore, model studies claim that more research is needed on benthic denitrification, since not well understood biogeochemical processes increase the uncertainties of future projections (Meier et al. 2011). Therefore another promising approach would be the use of a needle probe MIMS inlet to measure large sets of  $N_2/Ar$  profiles (Hilary and Seitzinger 2003) followed by modeling  $N_2$  production rates.

Both rivers, the Nemunas and the Oder, drain into lagoons before entering the Baltic Sea. Transformation processes within these lagoons have not been a main focus of this thesis but a recent study showed that the Szczecin lagoon act as an important nitrogen sink (Voss et al. 2010). However under hypoxia, which is increasing in the coastal zone of the Baltic Sea (Conley et al. 2011), nitrogen removal rates decrease with reactive ammonium accumulating through the process of dissimilatory reduction to ammonium (DNRA) (Jäntti et al. in press). Denitrification measurements performed during this thesis in muddy sediments of the Szczecin lagoon are the highest denitrification rates so far measured in the Baltic Sea and ranged from 5000 to 7500  $\mu\text{mol N m}^{-2} \text{d}^{-1}$  in summer (data not published). Simple budget calculations with the measured denitrification rates show that only 16 to 24% of the DIN load (46 kt DIN  $\text{yr}^{-1}$ , mainly in form of nitrate (Voss et al. 2010)) entering the lagoon from the Oder river is removed via denitrification. Therefore, still large amounts of nitrogen enter the adjacent coastal zone and these loads may even increase. Since the fate of nitrogen from rivers to the open Baltic Sea still is under debate, results from this thesis provide a valuable basis for future research to further study nitrogen transformation processes in estuaries and adjacent coastal areas to better understand how climate change and increasing direct anthropogenic pressures will influence the Baltic Sea. Projections of combined effects of climate change and eutrophication in general lead to the conclusion that if nothing is changed oxygen deficiencies and phytoplankton concentrations will continue to increase in the Baltic Sea (Meier et al. 2011). Combating eutrophication remains one of the major challenges in the Baltic Sea and worldwide.

## Bibliography

- Agedah, E.C., Binalaiyifa, H.E., Balli, A.S., Nedwell, D.B., 2009. Sources, turnover and bioavailability of dissolved organic nitrogen (DON) in the Colne estuary, UK. *Marine Ecology Progress Series* 23–33.
- Aleksandrov, S.V., 2010. Biological production and eutrophication of Baltic Sea estuarine ecosystems: The Curonian and Vistula Lagoons. *Marine Pollution Bulletin*, 61: 205–210.
- Algesten, G., Brydsten, L., Jonsson, P., Kortelainen, P., Lofgren, S., Rahm, L., Raike, A., Sobek, S., Tranvik, L., Wikner, J., and Jansson, M., 2006. Organic carbon budget for the Gulf of Bothnia: *J. Mar. Syst.*, 63, 155-161, 10.1016/j.jmarsys.2006.06.004.
- Alling, V., Humborg, C., Morth, C. M., Rahm, L., and Pollehne, F., 2008. Tracing terrestrial organic matter by delta S-34 and delta C-13 signatures in a subarctic estuary, *Limnol. Oceanogr.*, 53, 2594-2602.
- Alling, V., Sanchez-Garcia, L., Porcelli, D., Pugach, S., Vonk, J. E., van Dongen, B., Morth, C. M., Anderson, L. G., Sokolov, A., Andersson, P., Humborg, C., Semiletov, I., and Gustafsson, O., 2010. Nonconservative behavior of dissolved organic carbon across the Laptev and East Siberian seas, *Glob. Biogeochem. Cycle*, 24, 15, Gb403310.1029/2010gb003834.
- Altabet M. A. and W. G. Deuser, 1985. Seasonal variations in natural abundance of  $^{15}\text{N}$  in particles sinking to the deep Sargasso Sea. *Nature* 315: 218-219.
- Altabet, M. A. and J. J. McCarthy, 1985. Temporal and spatial variations in the natural abundance of  $^{15}\text{N}$  and PON from a warm-core ring. *Deep-Sea Research* 32(7): 755-772.
- Altabet, M. A. and J. J. McCarthy, 1986. Vertical patterns in  $^{15}\text{N}$  natural abundance in PON from the surface waters of warm-core rings. *Journal of Marine Research* 44: 185-201.
- Altabet, M. A. , 1988. Variations in nitrogen isotopic composition between sinking and suspended particles: implications for nitrogen cycling and particle transformation in the deep ocean. *Deep-Sea Research* 35(4): 535-554.
- Altabet, M. A. ,1989. A time-series study of the vertical structure of nitrogen and particle dynamics in the Sargasso Sea. *Limnology and Oceanography* 34(7): 1185-1201.
- Altabet, M.A., 2001. Nitrogen isotopic evidence for micronutrient control of fractional  $\text{NO}_3^-$  utilization in the equatorial Pacific. *Limnol. Oceanogr.*, 46(2): 368-380.
- Amon, R.M.W., Fitznar, H.-P., Benner, R., 2001. Linkages among the bioreactivity, chemical composition, and diagenetic state of marine dissolved organic matter. *Limnol. Oceanogr.*, 46, 2, 287–297.
- Andersen, J. H. et al., 2011. Getting the measure of eutrophication in the Baltic Sea: towards improved assessment principles and methods. *Biogeochemistry* 106, 137–156 DOI 10.1007/s10533-010-9508-4.
- Andrulewicz, E., 1997. An overview on lagoons in the Polish coastal area of the Baltic Sea. *International Journal of Salt Lake Research*, 6(2): 121-134.
- Antia, N. J., Harrison, P. J., Oliveira, L., 1991. The role of dissolved organic nitrogen in phytoplankton nutrition, cell biology and ecology. *Phycologia*, 30, 2, 1-89.
- BACC, 2008. Assessment of Climate Change for the Baltic Sea Basin. *Regional Climate Studies XII*. Springer Verlag, Berlin, 474 pp.
- Barlow R.G., Cummings D.G., Gibb S.W., 1997. Improved resolution of mono- and divinyl chlorophylls a and b and zeaxanthin and lutein in phytoplankton extracts using reverse phase C-8 HPLC. *Marine Ecology Progress Series* 161: 303-307.
- Behrendt, H., Dannowski, R., 2005. Nutrients and heavy metals in the Odra river system. *Weißensee Verlag Berlin*, 270 pp.
- Behrendt, H., Opitz, D., Kolanek, A., Korol, R., Stronska, M., 2008. Changes of the nutrient loads of the Odra River during the last century - their causes and consequences. *Journal of Water and Land Development* 12, 127-144.
- Benner, R., Biddanda, B., Black, B., and McCarthy, M., 1997. Abundance, size distribution, and stable

- carbon and nitrogen isotopic compositions of marine organic matter isolated by tangential-flow ultrafiltration, *Mar. Chem.*, 57, 243-263, 10.1016/S0304-4203(97)00013-3.
- Benner, R., 2002. Chemical Composition and Reactivity, in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Hansell, D. A., and Carlson, C. A., Academic Press, San Diego, 59-90.
- Berg, G.M., Repeta, D. J., and LaRoche, J., 2003b. The role of the picoeukaryote *Aureococcus anophagefferens* in cycling of marine high-molecular weight dissolved organic nitrogen, *Limnol. Oceanogr.*, 48, 1825-1830.
- Berg G.M., Glibert P.M., Jørgensen N.O.G., Balode M., Purina I., 2001. Variability in Inorganic and Organic Nitrogen Uptake Associated with Riverine Nutrient Input in the Gulf of Riga, Baltic Sea. *Estuaries* 24: 204-214.
- Berg G.M., Balode M., Purina I., Bekere S., Béchemin C., Maestrini S.Y., 2003a. Plankton community composition in relation to availability and uptake of oxidized and reduced nitrogen. *Aquatic Microbial Ecology* 30: 263-274.
- Bergström, S., Alexandersson, H., Carlsson, B., Josefsson, W., Karlsson, K-G., Westring, G., 2001. Climate and Hydrology of the Baltic Basin., in: *A Systems Analysis of the Baltic Sea*, edited by: Wulff, F., Rahm, L., and Larsson, P., Ecological Studies, Springer, Berlin.
- Berman T., 2001. The role of DON and the effect of N: P ratios on occurrence of cyanobacterial blooms: Implications from the outgrowth of *Aphanizomenon* in Lake Kinneret. *Limnology and Oceanography* 46: 443-447.
- Berman, T., 1997. Dissolved organic nitrogen utilization by an *Aphanizomenon* bloom in Lake Kinneret. *Journal of Plankton Research* 19, 577-586.
- Berman, T., and Bronk, D.A., 2003. Dissolved organic nitrogen: a dynamic participant in aquatic ecosystems: *Aquatic Microbial Ecology*, p. 279-305.
- Bernes, C., 2005. Change beneath the surface. Fälb & Hässler, 192 pp.
- Bitinas, A. et al., 2005. The results of integrated investigations of the Lithuanian coast of the Baltic Sea: geology, geomorphology, dynamics and human impact. *Geological Quarterly*, 49(4): 355-362.
- Bode, A., Varela, M. M., Teira, E., Fernandez, E., Gonzalez, N., Varela, M. M., 2004. Planktonic carbon and nitrogen cycling off northwest Spain: variations in production of particulate and dissolved organic pools. *Aquatic Microbial Ecology* 37, 1, 95-107.
- Böhlke, J.K., Mroczkowski, S.J. and Coplen, T.B., 2003. Oxygen isotopes in nitrate: new reference materials for  $18\text{O}:17\text{O}:16\text{O}$  measurements and observations on nitrate-water equilibration. *Rapid Communications in Mass Spectrometry*, 17(16): 1835-1846.
- Böttcher, J., Strebel, O., Voerkelius, S., and Schmidt, H.-L., 1990. Using isotope fractionation of nitrate-nitrogen and nitrate-oxygen for evaluation of microbial denitrification in a sandy aquifer: *Journal of Hydrology*, 114, p. 413-424.
- Boyer, E.W., Goodale, C.L., Jaworski, N.A., and Howarth, R.W., 2002. Anthropogenic nitrogen sources and relationships to riverine nitrogen export in the northeastern U.S.A.: *Biogeochemistry* 57/58, 137-169.
- Boyer, E.W., Howarth, R.W., Galloway, J.N., Dentener, F.J., Green, P.A., and Vörösmarty, C.J., 2006. Riverine nitrogen export from the continents to the coasts: *Global Biogeochemical Cycles*, 20, GB1S91;1-9.
- Bradley P.B., Sanderson M.P., Nejstgaard J.C., Sazhin A.F., Frischer M.E., Killberg-Thoreson L.M., Verity P.G., Campbell L., Bronk D.A., 2010. Nitrogen uptake by phytoplankton and bacteria during an induced *Phaeocystis pouchetii* bloom, measured using size fractionation and flow cytometric sorting. *Aquatic Microbial Ecology* 61:89-104.
- Brandes, J.A. and Devol, A.H., 1997. Isotopic fractionation of oxygen and nitrogen in coastal marine sediments. *Geochimica et Cosmochimica Acta*, 61(9): 1793-1801.
- Brandes, J.A. and Devol, A.H., 2002. A global marine-fixed nitrogen isotopic budget: Implications for Holocene nitrogen cycling. *Global Biogeochemical Cycles*, 16(4).



- Brandes, J.A., Devol, A.H., Yoshinari, T., Jayakumar, D.A. and Naqvi, S.W.A., 1998. Isotopic composition of nitrate in the central Arabian Sea and eastern tropical North Pacific: A tracer for mixing and nitrogen cycles. *Limnology and Oceanography*, 43(7): 1680-1689.
- Brandes, J. A., A. H. Devol, and C. Deutsch. 2007. New developments in the marine nitrogen cycle. *Chem. Rev.* 107, 577-589.
- Bronk D.A., Glibert P.M., 1993. Application of a  $^{15}\text{N}$  tracer method to the study of dissolved organic nitrogen uptake during spring and summer in Chesapeake Bay. *Marine Biology* 115: 501-508.
- Bronk, D.A., Glibert, P.M., Ward, B.B., 1994. Nitrogen Uptake, Dissolved Organic Nitrogen Release, and New Production. *Science* 265, 1843-1846.
- Bronk D.A., Ward B.B., 1999. Gross and net nitrogen uptake and DON release in the euphotic zone of Monterey Bay, California. *Limnology and Oceanography* 44 (3): 573-588.
- Bronk, D.A., Glibert, P.M., Malone, T.C., Banahan, S., and Sahlsten, E., 1998. Inorganic and organic nitrogen cycling in Chesapeake Bay: autotrophic versus heterotrophic processes and relationships to carbon flux: *Aquatic Microbial Ecology*, 15, 177-189.
- Bronk D.A., 2002. Dynamics of DON. In: Hansell DA, Carlson CA (eds) *Biogeochemistry of Marine Dissolved Organic Matter*. Elsevier, London, pp 153-200.
- Bronk D.A., See J.H., Bradley P., Killberg L., 2007. DON as a source of bioavailable nitrogen for phytoplankton. *Biogeosciences* 4: 283-296.
- Buchwald, C. and Casciotti, K.L., 2010. Oxygen isotopic fractionation and exchange during bacterial nitrite oxidation. *Limnology and Oceanography*, 55(3): 1064–1074.
- Burdige, D.J., 2002. Sediment pore waters. In: D.A. Hansell, C.A. Carlson (Eds.), *Biogeochemistry of Marine Dissolved Organic Matter*, Academic press, San Diego, 611-663.
- Burgman, J.O., Calles, B., Westman, F., 1987. Conclusions from a ten year study of oxygen-18 in precipitation and runoff in Sweden, In: Publications, IAEA Division of Publications (Eds.), *Isotope techniques in water resources development*. IAEA, 815pp.
- Codispoti, L. A., 2007. An oceanic fixed nitrogen sink exceeding  $400 \text{ Tg N a}^{-1}$  vs. the concept of homeostasis in the fixed-nitrogen inventory. *Biogeosciences*, 4, 233–253.
- Canfield, D.E., Glazer, A.N., and Falkowski, P.G., 2010, *The Evolution and Future of Earth's Nitrogen Cycle*: Science, 330, 192-196.
- Capone, D. G., J. P. Zehr, H. W. Paerl, B. Bergman, E. J. Carpenter, 1997. Trichodesmium, a globally significant marine cyanobacterium, *Science*, 276, 1221– 1229, 1997.
- Carstensen, J., D. J. Conley, J. H. Andersen, G. Aertebjerg, 2006. Coastal eutrophication and trend reversal: A Danish case study. *Limnol. Oceanogr.*, 51, 398-408.
- Casciotti, K.L., Sigman, D.M., Hastings, M.G., Böhlke, J.K. and Hilkert, A., 2002. Measurement of the Oxygen Isotopic Composition of Nitrate in Seawater and Freshwater Using the Denitrifier Method. *Anal. Chem.*, 74: 4905-4912.
- Casciotti, K. L., D. M. Sigman, and B. B. Ward, 2003. Linking diversity and stable isotope fractionation in ammonia-oxidizing bacteria. *Geomicrobiol. J.* 20: 335-353.
- Casciotti, K.L., McIlvin, M.R., 2007. Isotopic analyses of nitrate and nitrite from reference mixtures and application to Eastern Tropical North Pacific waters: *Marine Chemistry* 107, 184-201.
- Casciotti, K.L., McIlvin, M.R., Buchwald, C., 2010. Oxygen isotopic exchange and fractionation during bacterial ammonia oxidation. *Limnol. Oceanogr.*, 55, 2, 753–762.
- Casciotti, K. L. , Buchwald, C., Santoro, A. E., Frame, C., 2011. Assessment of Nitrogen and Oxygen Isotopic Fractionation During Nitrification and Its Expression in the Marine Environment. In artin G. Klotz, editor: *Methods in Enzymology*, Vol. 486, Burlington: Academic Press, pp. 253-280.
- Carlson, C. A., 2002. Production and Removal Processes, in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Hansell, D. A., and Carlson, C. A., Academic Press, San Diego, 91-151.
- Carlsson P., Graneli E., Tester P., Boni L., 1995. Influences of riverine humic substances on bacteria, protozoa, phytoplankton, and copepods in a coastal plankton community. *Marine Ecology Progress Series* 127: 13-221.



- Carlsson P., Graneli E., Segatto A.Z., 1999. Cycling of biologically available nitrogen in riverine humic substances between marine bacteria, a heterotrophic nanoflagellate and a photosynthetic dinoflagellate. *Aquatic Microbial Ecology* 18: 23-36.
- Cauwet, G., 2002. DOM in the coastal zone. In: Hansell, D. A., Carlson, C. A. (Eds.), *Biogeochemistry of Marine Dissolved Organic Matter*. Academic Press, San Diego, pp. 579–609.
- Christensen, J.H., Räisänen, J., Iversen, T., Bjørge, D., Christensen, O.B., Rummukainen, M., 2001. A synthesis of regional climate change simulations – a Scandinavian perspective. *Geophysical Research Letters* 28, 1003-1006.
- Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem. *Marine Ecology-Progress Series*, 210: 223-253.
- Conley, D.J., Björck, S., Bonsdorff, E., Carstensen, J., Destouni, G., Gustafsson, B.G., Hietanen, S., Kortekaas, M., Kuosa, H., Meier, H.E.M., Müller-Karulis, B., Nordberg, K., Norkko, A., Nürnberg, G., Pitkänen, H., Rabalais, N.N., Rosenberg, R., Savchuk, O.P., Slomp, C.P., Voss, M., Wulff, F., Zillén, L., 2009a. Hypoxia-Related Processes in the Baltic Sea. *Environmental Science & Technology* 43, 3412-3420.
- Conley, D.J., Humborg, C., Rahm, L., Savchuk, O.P., Wulff, F., 2002. Hypoxia in the Baltic Sea and Basin-Scale Changes in Phosphorous Biogeochemistry. *Environmental Science & Technology* 36, 5315-5320.
- Conley, D.J., Humborg, C., Smedberg, E., Rahm, L., Papush, L., Danielsson, Å., Clarke, A., Pastuszak, M., Aigars, J., Ciuffa, D., Mörtz, C.-M., 2008a. Past, present and future state of the biogeochemical Si cycle in the Baltic Sea. *Journal of Marine Systems* 73, 338-346.
- Conley, D.J., Humborg, C., Smedberg, E., Rahm, L., Papush, L., Danielsson, A., Clarke, A., Pastuszak, M., Aigars, J., Ciuffa, D., Morth, C.M., 2008b. Past, present and future state of the biogeochemical Si cycle in the Baltic Sea. *Journal of Marine Systems* 73, 338-346.
- Conley, D.J., Paerl, H.W., Howarth, R.W., Boesch, D.F., Seitzinger, S.P., Havens, K.E., Lancelot, C., Likens, G.E., 2009b. Controlling Eutrophication: Nitrogen and Phosphorus. *Science* 323, 1014-1015.
- Conley, D. J. et al., 2011. Hypoxia Is Increasing in the Coastal Zone of the Baltic Sea. *Environmental Science & Technology*, 45, 16, 6777-6783.
- Cornell, S., Rendell, A., and Jickells, T., 1995. Atmospheric inputs of dissolved organic nitrogen to the oceans. *Nature* 376, 243-246
- Costanza, R., et al., 1997. The value of the world's ecosystem services and natural capital. *Nature* 387, 253-260.
- Craig, H., and Gordon, L. I., 1965. Deuterium and oxygen 18 variations in the ocean and the marine atmosphere, in: *Stable Isotopes in Oceanographic Studies and Paleotemperatures*, edited by: Tongiorgi, E., Cons. Naz. Ric., Lab. Geol. Nucl., Pisa, 9-72.
- Dalsgaard, T., Canfield, D.E., Petersen, J., Thamdrup, B., Acuna-Gonzalez, J., 2003. N<sub>2</sub> production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica. *Nature* 422, 606–608.
- Dähnke, K., Serna, A., Blanz, T., Emeis, K.-C., 2008. Sub-recent nitrogen-isotope trends in sediments from Skagerrak (North Sea) and Kattegat: Changes in N-budgets and N-sources? *Marine Geology* 253, 92-98.
- Dähnke, K., Bahlmann, E. and Emeis, K., 2008. A nitrate sink in estuaries? An assessment by means of stable nitrate isotopes in the Elbe estuary. *Limnol. Oceanogr.*, 53(4): 1504-1511.
- Dähnke, K., Emeis, K., Johannsen, A. and Nagel, B., 2010. Stable isotope composition and turnover of nitrate in the German Bight. *Marine Ecology Progress Series*, 408: 7–18.
- Déqué, M., Rowell, D.P., Lüthi, D., Giorgi, F., Christensen, J.H., Rockel, B., Jacob, D., Kjellström, E., Castro, M.D., Hurk, B.V.D., 2007. An intercomparison of regional climate simulations for Europe: Assessing uncertainties in model projections. *Climatic Change* 81, 53-70.
- Deutsch, B., Forster, S., Wilhelm, M., Dippner, J.W., Voss, M., 2010. Denitrification in sediments as a major nitrogen sink in the Baltic Sea: an extrapolation using sediment characteristics. *Biogeosciences* 7, 3259–3271.

- Deutsch, B., Kahle, P. and Voss, M., 2006a. Assessing the source of nitrate pollution in water using stable N and O isotopes. *Agron. Sustain. Dev.*, 26: 263-267.
- Deutsch, B., Mewes, M., Liskow, I. and Voss, M., 2006b. Quantification of diffuse nitrate inputs into a small river system using stable isotopes of oxygen and nitrogen in nitrate. *Organic Geochemistry* 37: 1333-1342.
- Devol, A.H., 2003. Solution to a marine mystery. *Nature* 422, 575-576.
- Devol, A.H., 2008. Denitrification including Anammox. In: D. Capone, D. Bronk, M. Mulholland and E. Carpenter (Editors), *Nitrogen in the Marine Environment*. Academic Press: 263-301.
- Dippner, J.W., Ikaunieca, A., 2001. Long-term zoobenthos variability in the Gulf of Riga in relation to climate variability. *J. Mar. Syst.* 30, 155-164.
- Diaz, R.J. and Rosenberg, R., 2008. Spreading Dead Zones and Consequences for Marine Ecosystems. *Science* 321: 926-930.
- Dittmar, T., and Paeng, J., 2009. A heat-induced molecular signature in marine dissolved organic matter: *Nature Geoscience*, 2, 175-179, doi: 10.1038/ngeo440.
- Duce, R.A. et al., 2008. Impacts of atmospheric anthropogenic nitrogen on the open ocean. *Science* 320, 893-897.
- Dugdale, R.C. and Goering, J.J., 1967. Uptake of new and regenerated forms of nitrogen in primary productivity. *Limnol. Oceanogr.* 12: 169-206.
- Dugdale, R.C., and Wilkerson, F.P., 1986. The use of  $^{15}\text{N}$  to measure nitrogen uptake in eutrophic oceans; experimental considerations: *Limnology and Oceanography*, 31, 673-689.
- Dumont, E., Harrison, J.A., Kroeze, C., Bakker, E.J. and Seitzinger, S.P., 2005. Global distribution and sources of dissolved inorganic nitrogen export to the coastal zone: Results from a spatially explicit, global model. *Global Biogeochemical Cycles*, 19: 1-13.
- Elken, J., Matthäus, W., 2008. Assessment of Climate Change for the Baltic Sea Basin, In: team, BACC author team (Eds.), *Baltic Sea Oceanography*. Springer Verlag Berlin, Berlin, 379-385.
- Elmgren, R., 2001. Understanding Human Impact on the Baltic Ecosystem: Changing Views in Recent Decades. *Ambio* 30, 222-231.
- Elmgren, R., Larsson, U., 2001a. Eutrophication in the Baltic Sea area - Integrated Coastal Management Issues, In: Bodungen, B.v.Turner, R.K. (Eds.), *Science and Integrated Coastal Management*. Dahlem University Press, Berlin, pp. 15-35.
- Elmgren, R., Larsson, U., 2001b. Nitrogen and the Baltic Sea: Managing Nitrogen in Relation to Phosphorus, In: (Eds.), *2nd International Nitrogen Conference on Science and Policy*. The Scientific World, 371-377.
- Emeis, K., Christiansen, C., Edelvang, K., Jähmlich, S., Kozuch, J., Laima, M., Leipe, T., Löffler, A., Lund-Hansen, L.C., Miltner, A., Pazdro, K., Pempkowiak, J., Pollehne, F., Shimmield, T., Voss, M., Witt, G., 2002. Material Transport from the nearshore to the basinal environment in the Southern Baltic Sea II: Synthesis of data on origin and properties of material. *Journal of Marine Systems* 35, 151-168.
- Emeis, K., Struck, U., Leipe, T., Pollehne, F., Kunzendorf, H., Christiansen, C., 2000. Changes in the C, N, P burial rates in some Baltic Sea sediments over the last 150 years - relevance to P regeneration rates and the phosphorus cycle. *Marine Geology* 167, 43-59.
- Emery, K. O., 1968. Relict sands on continental shelves of the world. *AAPG Bull.* 52, 445-464.
- Engel, A., Goldthwait, S., Passow, U., Alldredge, A., 2002. Temporal decoupling of carbon and nitrogen dynamics in a mesocosm diatom bloom. *Limnology and Oceanography* 47, 753-761.
- Eriksson Hägg, H., C. Humborg, C.-M. Mörtz, M. Rodrigues Medina, F. Wulff, 2010. Scenario analysis on protein consumption and climate change effects on riverine N export to the Baltic Sea. *Environ. Sci. Technol.* 44: 2379-2385.
- Erisman, J.W., Sutton, M.A., Galloway, J., Klimont, Z., and Winiwarter, W., 2008. How a century of ammonia synthesis changed the world: *nature geoscience*, v. 1, p. 636- 639.
- Feistel, R., G. Nausch and N. Wasmund, Eds., 2008. *State and evolution of the Baltic Sea, 1952 – 2005. A detailed 50-year survey of meteorology and climate, physics, chemistry, biology, and marine environment*. Hoboken: Wiley-Interscience. 703 S.

- Finni, T., Kononen, K., Olsonen, R. and Wallström, K., 2001. The history of cyanobacterial blooms. *Ambio*, 30(4-5): 172-178.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K. G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N., 2006. Climate-carbon cycle feedback analysis: Results from the (CMIP)-M-4 model intercomparison, *Journal of Climate*, 19, 3337-3353.
- Fisher, D.C., Oppenheimer, M., 1991. Atmospheric Nitrogen Deposition and the Chesapeake Bay Estuary. *Ambio* 3/4, 102-108
- Fonselius, S., 2003. One hundred years of hydrographic measurements in the Baltic Sea. *Journal of Sea Research* 49, 229– 241.
- Fonselius, S.H., 1969. Hydrography of the Baltic Deep Basins. Fish Board of Sweden, Ser. Hydrogr. 97pp.
- Freeman, C., Evans, C.D., Monteith, D.T., Reynolds, B., Fenner, N., 2001a. Export of organic carbon from peat soils. *Nature* 412, 785-785.
- Freeman, C., Ostle, N., Kang, H., 2001b. An enzymic 'latch' on a global carbon store - A shortage of oxygen locks up carbon in peatlands by restraining a single enzyme. *Nature* 409, 149-149.
- Fry, B., 2002. Conservative mixing of stable isotopes across estuarine salinity gradients: A conceptual framework for monitoring watershed influences on downstream fisheries production. *Estuaries*, 25(2): 264-271.
- Fry, B., 2006. *Stable Isotope Ecology*. Springer, New York, NY.
- Fry, B. and Sherr, E.B., 1984.  $\delta^{13}\text{C}$  measurements as indicators of carbon flow in marine and freshwater ecosystems. *Contributions to Marine Science* 27: 13-47.
- Fröhlich, K., Grabczak, J., and Rozanski, K., 1988. Deuterium and Oxygen-18 in the Baltic Sea, *Chemical Geology*, 72, 77-83.
- Fuhrman, J.A., 1999. Marine viruses and their biogeochemical and ecological effects. *Nature* 399:541-548.
- Gao, H. et al., 2010. Aerobic denitrification in permeable Wadden Sea sediments. *Isme Journal*, 5(4): 776-776.
- Gao, H. et al., 2012. Intensive and extensive nitrogen loss from intertidal permeable sediments of the Wadden Sea. *Limnol. Oceanogr.*, 57, 1, 185–198.
- Galloway, J.N., Aber, J.D., Erisman, J.W., Seitzinger, S.P., Howarth, R.W., Cowling, E.B., and Cosby, B.J., 2003. The Nitrogen Cascade: *BioScience*, 53, 341-356.
- Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer, E.W., Howarth, R.W., Seitzinger, S.P., Asner, G.P., Cleveland, C.C., Green, P.A., Holland, E.A., Karl, D.M., Michaels, A.F., Porter, J.H., Townsend, A.R., and Vörösmarty, C.J., 2004. Nitrogen cycles: past, present, and future: *Biogeochemistry*, 70, 153-226.
- Gasiunaite, Z.R., Daunys, D., Olenin, S. and Razinkovas, A., 2008. The Curonian Lagoon. In: M.M. Caldwell et al. (Editors), *Ecology of Baltic Coastal Waters*. Ecological Studies. Springer Berlin Heidelberg, pp. 197-215.
- Gasol, J.M., Casamayor, E.O., Joint, I., Garde, K., Gustavson, K., Benlloch, S., Díez, B., Schauer, M., Massana, R., Pedrós-Alió, C., 2004. Control of heterotrophic prokaryotic abundance and growth rate in hypersaline planktonic environments. *Aquatic Microbiological Ecology* 34: 193–206.
- Gerlach, S.A., 1994. Oxygen conditions improve when the salinity in the Baltic Sea decreases. *Marine Pollution Bulletin* 28, 413-416.
- Glibert, P.M., Bronk, D.A., 1994. Release of Dissolved Organic Nitrogen by Marine diazotrophic Cyanobacteria, *Trichodesmium* spp.. *Applied and Environmental Microbiology* 60: 3996-4000.

- Giesler, R., Högberg, M.N., Strobel, B.W., Richter, A., Nordgren, A., Högberg, P., 2007. Production of dissolved organic carbon and low-molecular weight organic acids in soil solution driven by recent tree photosynthate. *Biogeochemistry* 84, 1-12.
- Giorgi, F., Hewitson, B., Christensen, J., Hulme, M., von Storch, H., Whetton, P., Jones, R., Mearns, L., Fu, C., 2001. Regional climate information – evaluation and projections, In: 2001, I.C.C. (Eds.), *The scientific basis contribution of working group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge New York, pp. 583-638.
- Graham, L.P., 2004. Climate change effects on river flow to the Baltic Sea. *Ambio* 33, 235-241.
- Graham, L.P., Hagemann, S., Jaun, S., Beniston, M., 2007. On interpreting hydrological change from regional climate models. *Climatic Change* 81, 97-122.
- Graneli, E., Wallstroem, K., Larsson, U., Graneli, W., Elmgren, R., 1990. Nutrient limitation of primary production in the Baltic Sea area. *Ambio* 19, 142-151.
- Granger, J., Sigman, D.M., Needoba, J.A. and Harrison, P.J., 2004. Coupled nitrogen and oxygen isotope fractionation of nitrate during assimilation by cultures of marine phytoplankton. *Limnol. Oceanogr.*, 49(5): 1763-1773.
- Granger, J., Sigman, D.M., Lehmann, M.F., and Tortell, P.D., 2008. Nitrogen and oxygen isotope fractionation during dissimilatory nitrate reduction by denitrifying bacteria: *Limnology and Oceanography*, 53, 2533–2545.
- Granger, J., Sigman, D.M., Rohde, M.M., Maldonado, M.T. and Tortell, P.D., 2010. N and O isotope effects during nitrate assimilation by unicellular prokaryotic and eukaryotic plankton cultures. *Geochimica et Cosmochimica Acta*, 74: 1030–1040.
- Grasshoff, K., Ehrhardt, M. and Kremling, K., 1983. *Methods of Seawater Analysis*. Verlag Chemie.
- Grelowski, A., Pastuszak, M., Sitek, S. and Witek, Z., 2000. Budget calculations of nitrogen, phosphorus and BOD5 passing through the Oder estuary. *Journal of Marine Systems* 25: 221-237.
- Gruber, N. and Galloway, J.N., 2008. An Earth-system perspective of the global nitrogen cycle. *nature*, 451: 293-296.
- Guo, L. D., Tanaka, N., Schell, D. M., and Santschi, P. H., 2003. Nitrogen and carbon isotopic composition of high-molecular-weight dissolved organic matter in marine environments, *Mar. Ecol.-Prog. Ser.*, 252, 51-60.
- Guo, L. D., White, D. M., Xu, C., and Santschi, P. H., 2009. Chemical and isotopic composition of high-molecular-weight dissolved organic matter from the Mississippi River plume, *Mar. Chem.*, 114, 63-71, 10.1016/j.marchem.2009.04.002.
- Gurrath, P., 2009. *Landwirtschaft in Deutschland und der Europäischen Union 2009*. Statistisches Bundesamt, Wiesbaden.
- Gustafsson, M., K. Nordberg, 2000. Living (Stained) Benthic Foraminifera and their Response to the Seasonal Hydrographic Cycle, Periodic Hypoxia and to Primary Production in Havstens Fjord on the Swedish West Coast. *Estuarine, Coastal and Shelf Science* 51, 743–761.
- Hagemann, S., Jacob, D., 2007. Gradient in the climate change signal of European discharge predicted by a multi-model ensemble. *Climatic Change* 81, 309-327.
- Hagen, E., Feistel, R., 2005. Climatic turning points and regime shifts in the Baltic Sea region: the Baltic winter index (WIBIX) 1659-2002. *Boreal Environment Research* 10, 211-224.
- Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., and Watson, R., 2008. A Global Map of Human Impact on Marine Ecosystems: *Science* 319, 948-952.
- Hannig, M., Lavik, G., Kuypers, M.M.M., Woebken, D., Martens-Habbena, W., Jürgens, K., 2007. Shift from denitrification to anammox after inflow events in the central Baltic Sea. *Limnology and Oceanography* 52, 1336–1345.

- Hedges, J.I. et al., 1986. Composition and fluxes of particulate organic material in the Amazon River. *Limnology and Oceanography*, 31: 717- 738.
- Hedges, J. I., Keil, R. G., Benner, R., 1997. What happens to terrestrial organic matter in the ocean? *Organic Geochemistry* 27, 5-6, 195–212.
- Hedges, J. I., 2002. Why dissolved organic matter? In: Hansell, D. A., Carlson, C. A. (Eds.), *Biogeochemistry of Marine Dissolved Organic Matter*. Academic Press, San Diego, pp. 1– 33.
- Heip, C.H.R. et al., 1995. Production and consumption of biological particles in temperate tidal estuaries, *Oceanography and marine Biology*, 33: 1-149.
- HELCOM, 2004. Baltic Sea Environment Proceedings: The Fourth Baltic Sea Pollution Load Compilation. Baltic Marine Environment Protection Commission, Helsinki, 188 pp.
- HELCOM, 2005. Nutrient Pollution to the Baltic Sea in 2000 Baltic Sea Environment Proceedings 100, [http://www.helcom.fi/publications/bsep/en\\_GB/bseplist/](http://www.helcom.fi/publications/bsep/en_GB/bseplist/).
- HELCOM, 2007. Baltic Sea Action Plan. Baltic Sea Environment Proceedings [http://www.helcom.fi/BSAP/ActionPlan/en\\_GB/ActionPlan/](http://www.helcom.fi/BSAP/ActionPlan/en_GB/ActionPlan/).
- HELCOM, Laane, A., Wilhelms, S., 1993. Second Baltic Sea Pollution load compilation. *Baltic Sea Environment Proceedings* 45, 1-162.
- HELCOM, 1996. Baltic Sea Environment Proceedings: Third periodic assessment of the state of the marine environment of the Baltic Sea. *Baltic Sea Environment Proceedings* No. 64B. Baltic Marine Environment Protection Commission, Helsinki, pp.1-252.
- HELCOM, 2009. Eutrophication in the Baltic Sea - An integrated thematic assessment of the effects of nutrient enrichment in the Baltic Sea region. *Baltic Sea Environment Proceedings* No. 115B. Baltic Marine Environment Protection Commission, Helsinki, pp. 1-152.
- Herlemann, D.P.R., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J.J., Andersson, A.F., 2011. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME Journal*: in press.
- Hertel, O., Skjøth, C.A., Brandt, J., Christensen, J.H., Frohn, L.M., Frydendall, J., 2003. Operational mapping of atmospheric nitrogen deposition to the Baltic Sea. *Atmosphere Chemistry and Physics* 3, 2083-2099.
- Heyen, H., Zorita, E., Storch, H.v., 1996. Statistical downscaling of monthly mean North Atlantic air pressure to sea level anomalies in the Baltic Sea. *Tellus* 48A, 312-323.
- Hietanen, S., Kuparinen, J., 2008. Seasonal and short-term variation in denitrification and anammox at a coastal station on the Gulf of Finland, Baltic Sea. *Hydrobiologia* 596:, 67-77, DOI 10.1007/s10750-10007-19058-10755.
- Hietanen, S., Lukkari, K., 2007. Effects of short-term anoxia on benthic denitrification, nutrient fluxes and phosphorus forms in coastal Baltic sediment. *Aquatic Microbial Ecology* 49, 293-302, doi: 210.3354/ame01146.
- Hilairy, E. H., Seitzinger. S. P., 2003. High-Resolution nitrogen gas profiles in sediment porewaters using a new membrane probe for membrane-inlet mass spectrometry (MIMS). *Marine Chemistry*, 83, 23–30.
- Hood, E., Scott, D., 2008. Riverine organic matter and nutrients in southeast Alaska affected by glacial coverage. *Nature Geoscience* 1, 583-587, doi:510.1038/ngeo1280.
- Howarth, R. W., Anderson, D., Cloern, J., Elfring, C., Hopkinson, C., Lapointe, B., Malone, T., Marcus, N., McGlathery, K., Sharpley, A., Walker, D., 2000. Nutrient pollution of coastal rivers, bays and seas. *Issues in Ecology* 7. Ecological Society of America, Washington DC.
- Howarth, R.W., Marino, R., 2006. Nitrogen as the limiting nutrient for eutrophication in coastal marine ecosystems: Evolving views over three decades. *Limnology and Oceanography* 51, 1, 2, 364–376.
- Howarth, R.W., 2008. Coastal nitrogen pollution: A review of sources and trends globally and regionally: *Harmful Algae* 8, 14-20.
- Huettel, M. and Gust, G., 1992. Impact of bioturbation on interfacial solute exchange in permeable sediments. *Marine ecology progress series*, 89(2-3): 253-267.

- Humborg, C., Morth, C.M., Sundbom, M., Wulff, F., 2007. Riverine transport of biogenic elements to the Baltic Sea - past and possible future perspectives. *Hydrology and Earth System Sciences* 11, 1593-1607.
- Humborg, C., Rahm, L., Conley, D.J., Tamminen, T., von Bodungen, B., 2008. Silicon and the Baltic Sea Long-term Si decrease in the Baltic Sea - A conceivable ecological risk? *Journal of Marine Systems* 73, 221-222.
- Humborg, C., Morth, C. M., Sundbom, M., Borg, H., Blenckner, T., Giesler, R., and Ittekkot, V., 2010. CO<sub>2</sub> supersaturation along the aquatic conduit in Swedish watersheds as constrained by terrestrial respiration, aquatic respiration and weathering, *Glob. Change Biol.*, 16, 1966-1978, 10.1111/j.1365-2486.2009.02092.x.
- Humborg, C., Smedberg, E., Medina, M. R., and Morth, C. M., 2008. Changes in dissolved silicate loads to the Baltic Sea - The effects of lakes and reservoirs, *J. Mar. Syst.*, 73, 223-235, 10.1016/j.jmarsys.2007.10.014.
- Hunter, E.M., Mills, H.J. and Kostka, J.E., 2006. Microbial community diversity associated with carbon and nitrogen cycling in permeable shelf sediments. *Applied and Environmental Microbiology*, 72(9): 5689-5701.
- Hurrell, J.W., 1995. Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and Precipitation. *Science* 269, 676-679.
- IPCC: Climate Change, 2001. The Scientific Basis - Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, New York, pp 881.
- Jahnke, R. et al., 2005. Organic matter remineralization and porewater exchange rates in permeable South Atlantic Bight continental shelf sediments. *Continental Shelf Research*, 25(12-13): 1433-1452.
- Janssen, F., Huettel, M. and Witte, U., 2005. Pore-water advection and solute fluxes in permeable marine sediments (II): Benthic respiration at three sandy sites with different permeabilities (German Bight, North Sea). *Limnology and Oceanography*, 50(3): 779-792.
- Jäntti, H., Stange, F., Leskinen, E. and Hietanen, S., 2011. Seasonal variation in nitrification and nitrate-reduction pathways in coastal sediments in the Gulf of Finland, Baltic Sea. *Aquatic Microbial Ecology*, 63(2): 171-181.
- Jäntti, H., Hietanen, S., 2011. The effects of hypoxia on sediment nitrogen cycling in the Baltic Sea. *AMBIO*, in press.
- Jeffrey, S.W. and Welschmeyer, N.A., 1997. Spectrophotometric and fluorometric equations in common use in oceanography. In: S.W. Jeffrey, R.F.C. Mantoura and S.W. Wright (Editors), *Phytoplankton pigments in oceanography: guidelines to modern method*. UNESCO Publishing, Paris, pp. 597-615.
- Jensen, M.M., Lam, P., Revsbech, N.P., Nagel, B., Gaye, B., Jetten, M.S.M., Kuypers, M.M.M., 2011. Intensive nitrogen loss over the Omani Shelf due to anammox coupled with dissimilatory nitrite reduction to ammonium. *ISME Journal* 5, 1660-1670.
- Jones, M.N., 1984. Nitrate reduction by shaking with cadmium: Alternative to cadmium columns. *Water Research* 18: 643-646.
- Jonsson, P., Carman, R., 1994. Changes in deposition of organic matter and nutrients the Baltic Sea during the twentieth century. *Marine Pollution Bulletin* 28, 417-426.
- Jorgensen, N.G., Tranvik, L.J., Berg, G.M., 1999. Occurrence and bacterial cycling of dissolved nitrogen in the Gulf of Riga, the Baltic Sea. *Marine Ecology Progress Series* 191, 1-18.
- Kaldy, J. E., Cifuentes, L. A., and Brock, D., 2005. Using stable isotope analyses to assess carbon dynamics in a shallow subtropical estuary, *Estuaries*, 28, 86-95, 10.1007/bf02732756.
- Kendall, C., 1998. Tracing nitrogen sources and cycling in catchments. In: C. Kendall and J.J. McDonnell (Editors), *Isotope tracers in catchment hydrology*. Elsevier, Amsterdam, pp. 521-576.

- Kendall, C., Elliott, E.M., and Wankel, S.D., 2007. Tracing anthropogenic inputs of nitrogen to ecosystems, Chapter 12, In: R.H. Michener and K. Lajtha (Eds.), *Stable Isotopes in Ecology and Environmental Science*, 2nd edition, Blackwell Publishing, p. 375- 449.
- Kester, D.R., Duedall, I.W., Connors, D.N., Pytkowic, R.M., 1967. Preparation of artificial Seawater. *Limnology and Oceanography* 12: 176-179.
- Koch, B.P., Ludwichowski, K.-U., Kattner, G., Dittmar, T., and Witt, M., 2008. Advanced characterization of marine dissolved organic matter by combining reversed-phase liquid chromatography and FT-ICR-MS: *Marine Chemistry* 111, 233-241.
- Knapp, A.N., D, M. Sigman, F. Lipschultz. 2005. N isotopic composition of dissolved organic nitrogen and nitrate at the Bermuda Atlantic Time-series Study site. *GLOBAL BIOGEOCHEMICAL CYCLES*, VOL. 19, GB1018, doi:10.1029/2004GB002320.
- Knapp, A.N., D, M. Sigman, F. Lipschultz, A. B. Kustka, D. G. Capone, 2011. Interbasin isotopic correspondence between upper-ocean bulk DON and subsurface nitrate and its implications for marine nitrogen cycling. *GLOBAL BIOGEOCHEMICAL CYCLES*, VOL. 25, GB4004, doi:10.1029/2010GB003878.
- Knowles, R., 1982. Denitrification. *Microbial. Rev.* 46, 43-70.
- Köppen, W., Geiger, R., 1960. *Karten der Klimatypen - Climate Maps*. Brockhaus Verlag, Wiesbaden, pp.
- Koroleff, F., 1983. Determination of total phosphorus. In: Grasshoff, K., Erhardt, M., Kremling, K. (eds) *Methods for Seawater Analysis*. Verlag Chemie, Weinheim pp 167-173.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift* 15, 259-263.
- Korth, F., Deutsch, B., Liskow, I. and Voss, M., 2011. Uptake of dissolved organic nitrogen by size-fractionated plankton along a salinity gradient from the North Sea to the Baltic Sea. *Biogeochemistry*: 1-14. DOI 10.1007/s10533-011-9656-1.
- Krämer, I. Hürdler, J., Hirschfeld, J., Vehnohr, M., Schernewski, G., 2011. Nutrient Fluxes from Land to Sea: Consequences of Future Scenarios on the Oder River Basin – Lagoon – Coastal Sea System. *Internat. Rev. Hydrobiol.* 96, 5, 520–540.
- Kullman, L., 2007. Tree line population monitoring of *Pinus sylvestris* in the Swedish Scandes, 1973-2005: implications for tree line theory and climate change ecology. *Journal of Ecology* 95, 41-52.
- Kullman, L., Kjallgren, L., 2006. Holocene pine tree-line evolution in the Swedish Scandes: Recent tree-line rise and climate change in a long-term perspective. *Boreas* 35, 159-168.
- Kuypers, M.M.M., Sliekers, A.O., Lavik, G., Schmid, M., Jorgensen, B.B., Kuenen, J.G., Damste, J.S.S., Strous, M., Jetten, M.S.M., 2003. Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. *Nature* 422, 608–611.
- Lääne, A., Kraav, E., Titova, G., 2005. Global International Waters Assessment, Baltic Sea, GIWA Regional assessment 17, p 1-69.
- Larsson, U., Elmgren, R., Wulff, F., 1985. Eutrophication and the Baltic Sea: causes and consequences. *Ambio* 14, 9-14.
- Larsson, J., Gustafsson, O., and Ingri, J., 2002. Evaluation and optimization of two complementary cross-flow ultrafiltration systems toward isolation of coastal surface water colloids, *Environmental Science & Technology*, 36, 2236-2241, 10.1021/es010325v.
- Lass, H.U., Mohrholz, V. and Seifert, T., 2001. On the dynamics of the Pomeranian Bight. *Continental Shelf Research*, 21(11-12): 1237-1261.
- Lass, H. U., Mohrholz, V., and Seifert, T., 2005. On pathways and residence time of saltwater plumes in the Arkona Sea, *J. Geophys. Res.-Oceans*, 110, 24, C1101910.1029/2004jc002848.
- Laudon, H., Sjöblom, V., Buffam, I., Seibert, J., Morth, M., 2007. The role of catchment scale and landscape characteristics for runoff generation of boreal streams. *Journal of Hydrology* 344, 198-209.
- Lee, S., Fuhrman, J.A., 1987. Relationship between Biovolume and Biomass of Naturally

- Derived Marine Bacterioplankton. *Applied and Environmental Microbiology* 53: 1298-1303.
- Lehmann, M.F., Reichert, P., Bernasconi, S.M., Barbieri, A., and McKenzie, J.A., 2003. Modelling nitrogen and oxygen isotope fractionation during denitrification in a lacustrine redox-transition zone: *Geochimica et Cosmochimica Acta*, 67, 2529-2542.
- Lehmann, M.F., Sigman, D.M. and Berelson, W.M., 2004. Coupling the  $15\text{N}/14\text{N}$  and  $18\text{O}/16\text{O}$  of nitrate as a constraint on benthic nitrogen cycling. *Marine Chemistry*, 88: 1-20.
- Lehmann, M.F. et al., 2007. The distribution of nitrate  $15\text{N}/14\text{N}$  in marine sediments and the impact of benthic nitrogen loss on the isotopic composition of oceanic nitrate. *Geochimica et Cosmochimica Acta*, 71(22): 5384-5404.
- Letscher, R. T., Hansell, D. A., and Kadko, D., 2011. Rapid removal of terrigenous dissolved organic carbon over the Eurasian shelves of the Arctic Ocean, *Mar. Chem.*, 123, 78-87, 10.1016/j.marchem.2010.10.002.
- Lindroth, P., Mopper, K., 1997. High Performance Liquid Chromatographic Determination of Subpicomole Amounts of Amino Acids by Precolumn Fluorescence Derivatization with o-Phthaldialdehyde. *Analytical Chemistry* 51: 1667-1674.
- Liu, K.-K., and Kaplan, I.R., 1989, The eastern tropical Pacific as a source of  $15\text{N}$ -enriched nitrate in seawater off southern California: *Limnol. Oceanogr.*, 34, 820-830.
- Løvdaal, T., Eichner T., Grossart H.-P., Carbone V., Chou L., Thingstad T.F., 2008. Competition for inorganic and organic forms of nitrogen and phosphorous between phytoplankton and bacteria during an *Emiliania huxleyi* spring bloom (PeECE II). *Biogeosciences* 5: 371–383.
- Macko, S. A., and Estep, M. L. F.: Microbial alteration of stable nitrogen and carbon isotopic compositions of organic matter, *Org. Geochem.*, 6, 787-790, 1984.
- Mariotti, A., Landreau, A., and Simon, B., 1988.  $15\text{N}$  isotope biogeochemistry and natural denitrification process in groundwater: Application to the chalk aquifer of northern France: *Geochimica et Cosmochimica Acta* 52, 1869-1878.
- Matthäus, W., Franck, H., 1992. Characteristics of major Baltic inflows- a statistical analysis. *Continental Shelf Research* 12, 1375-1400.
- Matthäus, W., Schinke, H., 1999. The influence of river runoff on deep water conditions of the Baltic Sea. *Hydrobiologica* 393, 1-10.
- Mayer, B., S. M. Bollwerk, T. Mansfeldt, B. Hutter, and J. Veizer. 2001. The oxygen isotope composition of nitrate generated by nitrification in acid forest floors. *Geochim. Cosmochim. Acta* 65, 2743-2756.
- Mayer, B., Boyer, E.W., Goodale, C., Jaworski, N.A., Breemen, N.v., Howarth, R.W., Seitzinger, S., Billen, G., Lajtha, K., Nadelhoffer, K., Dam, D.v., Hetling, L.J., Nosal, M., and Paustian, K., 2002. Sources of nitrate in rivers draining sixteen watersheds in the northeastern U.S.: Isotopic constraints: *Biogeochemistry*, 57/58, 171-197.
- McCallister, S. L., Bauer, J. E., Ducklow, H. W., and Canuel, E. A., 2006. Sources of estuarine dissolved and particulate organic matter: A multi-tracer approach, *Organic Geochemistry*, 37, 454-468, 10.1016/j.orggeochem.2005.12.005.
- McCallister, S.L., Bauer, J.E., Canuel, E.A., 2006. Bioreactivity of estuarine dissolved organic matter: A combined geochemical and microbiological approach. *Limnology and Oceanography* 51: 94–100.
- McClelland, J.W. and Valiela, I., 1998. Linking nitrogen in estuarine producers to land-derived sources. *Limnol. Oceanogr.*, 43(4): 577-585.
- McClelland, J.W., Valiela, I. and Michener, R.H., 1997. Nitrogen-stable isotope signatures in estuarine food webs : A record of increasing urbanization in coastal watersheds. *Limnol. Oceanogr.*, 42: 930-937.
- Meier, H.E.M., 2006. Baltic Sea climate in the last twenty-first century: A dynamical downscaling approach using two global models and two emission scenarios. *Climate Dynamics* 27, 39-68 doi: 101007/s100382-101006-100124-x.



- Meier, H.E.M., Kauker, F., 2003. Modelling decadal variability of the Baltic Sea. Part II: Role of freshwater inflow and large-scale atmospheric circulation for salinity. *Journal of Geophysical Research* 108 3368 doi:10.1029/102003JC001799.
- Meier, H.E.M., Eilola, K., Almroth, E., 2011. Climate-related changes in marine ecosystems simulated with a 3-dimensional coupled physical-biogeochemical model of the Baltic Sea. *Climate Research*, 48, 31–55.
- Meybeck, M., 1988. How to establish and use world budgets of riverine material, in: *Physical and chemical weathering in geochemical cycles*, edited by: Lerman, A., and Meybeck, M., Kluwer Academic, Dordrecht, 247-272.
- Middelburg, J.J. and Nieuwenhuize, J., 1998. Carbon and nitrogen stable isotopes in suspended matter and sediments from the Schelde Estuary. *Marine Chemistry*, 60(3-4): 217-225.
- Middelburg, J.J. and Nieuwenhuize, J., 2000. Uptake of dissolved inorganic nitrogen in turbid, tidal estuaries. *Marine Ecology Progress Series*, 192: 79-88.
- Middelburg, J.J. and Nieuwenhuize, J., 2001. Nitrogen isotope tracing of dissolved inorganic nitrogen behaviour in tidal estuaries. *Estuar. Coast. Shelf Sci.* 53, 385-391.
- Middelburg, J.J., Soetaert, K., Herman, P.M.J. and Heip, C.H.R., 1996. Denitrification in marine sediments: A model study. *Global Biogeochem. Cycles*, 10(4): 661-673.
- Mingawa, M., Wada, E., 1986. Nitrogen isotope ratios of red tide organisms in the East China Sea: A characterization of biological nitrogen fixation. *Mar Chem* 19:245-256.
- Mohrholz, V., Pastuszak, M., Sitek, S.a., Nagel, K. and Lass, H., 1998. The exceptional Oder flood in summer 1997- riverine mass and nutrient transport into the Pomeranian Bight. *Ocean Dynamics*, 50(2): 129-144.
- Montoya, J.P., 20. Nitrogen Stable Isotopes in Marine Environments. In Capone, D. G., D. Bronk, M. Mulholland, and E. J. Carpenter [eds.]. *Nitrogen in the marine environment*, 2nd edition. Elsevier Science, pp. 1277-1297.
- Moran, M. A., and Zepp, R. G.: Role of photoreactions in the formation of biologically labile compounds from dissolved organic matter, *Limnol. Oceanogr.*, 42, 1307-1316, 1997.
- Mort, H.P., Slomp, C.P., Gustafsson, B.G., and Andersen, T.r.J., 2010. Phosphorus recycling and burial in Baltic Sea sediments with contrasting redox conditions: *Geochimica et Cosmochimica Acta*, 74, 1350-1362.
- Mudelsee, M., Börngen, M., Tetzlaff, G., Grünwald, U., 2004. Extreme floods in central Europe over the past 500 years: Role of cyclone pathway "Zugstrasse Vb". *Journal of Geophysical Research* 109, No. D23101, doi:23110.21029/22004JD005034.
- Mulholland M.R., Glibert P.M., Berg G.M., Van Heukelem L., Pantoja S., Lee C., 1998. Extracellular amino acid oxidation by phytoplankton and cyanobacteria: a cross-ecosystem comparison. *Aquatic microbial ecology* 15:141-152.
- Mulholland M.R., Bronk A.B., Capone D.G., 2004. Dinitrogen fixation and release of ammonium and dissolved organic nitrogen by *Trichodesmium* IMS101. *Aquatic Microbial Ecology* 37: 85-94.
- Mulholland M.R., Lomas M.W., 2008. Nitrogen Uptake and Assimilation In: Capone D.G., Bronk D.A., Mulholland M.R., Carpenter E.J. (eds) *Nitrogen in the marine environment*. Elsevier, New York, pp 303-384.
- Nausch, G., Nehring, D., Nagel, K., 2008. Nutrient concentrations, trends and their relation to eutrophication. In: Feistel R, Nausch G, Wasmund N (eds) *State and Evolution of the Baltic Sea, 1952-2005*. Wiley, New Jersey, pp 337-366.
- Nausch, G., Feistel, R., Umlauf, L., Nagel, K., and Siegel, H., 2010. *Hydrographisch-chemische Zustandseinschätzung der Ostsee*, Leibniz- Institut für Ostseeforschung Warnemünde, Warnemuende.
- Neumann, G. H., Fonselius, S., and Wahlman, L., 1959. Measurements on the content of non-volatile organic material in atmospheric precipitation, *International Journal of Air Pollution*, 2, 132-141.

- Neumann, T., 2000. Towards a 3D-ecosystem model of the Baltic Sea. *Journal Marine Systems* 25, 405-420.
- Neumann, T., Fennel, W., Kremp, C., 2002. Experimental Simulations with an Ecosystem Model of the Baltic Sea: A Nutrient Load Reduction Experiment. *Global Biogeochemical Cycles* 16, 7-19.
- Neumann, T., 2007. The fate of river-borne nitrogen in the Baltic Sea- An example for the River Oder. *Estuarine, Coastal and Shelf Science* 73: 1-7.
- Nielsen, L. P., 1992. Denitrification in sediment determined from nitrogen isotope pairing. *FEMS Microbiology Letters*, 86, 357-362.
- Nixon, S.W., 1995. Coastal marine eutrophication: A definition, social causes, and future concerns: *Ophelia*, 49, 199-219.
- Norrman, B., Zweifel, U. L., Hopkinson, C. S., and Fry, B., 1995. Production and utilization of dissolved organic-carbon during an experimental diatom bloom. *Limnol. Oceanogr.*, 40, 898-907.
- Ogawa, H., Amagai, Y., Koike, I., Keiser, K., Benner, R., 2001. Production of refractory dissolved organic matter by bacteria. *Science* 292, 917-920.
- Ohlendorf, U., Stühr, A., Siegmund, H., 2000. Nitrogen fixation by diazotrophic cyanobacteria in the Baltic Sea and transfer of the newly fixed nitrogen to picoplankton organisms. *Journal of Marine Research* 25: 213-220.
- Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., 2007. *Vegan: Community Ecology Package*. R package version 1.8-7. <http://r-forge.rproject.org/projects/vegan/>.
- Omstedt, A., Elken, J., Lehmann, A., Piechura, J., 2004. Knowledge of the Baltic Sea physics gained during the BALTEX and related programmes. *Progress in Oceanography* 63, 1-28.
- Omstedt, A., Gustafsson, B., Rohde, J., Walin, G., 2000. Use of Baltic Sea modelling to investigate the water cycle and heat balance in GCM and regional climate models. *Climate Research* 15, 95-108.
- Otero, E., Culp, R., Noakes, J. E., and Hodson, R. E., 2003. The distribution and delta C-13 of dissolved organic carbon and its humic fraction in estuaries of southeastern USA, *Estuar. Coast. Shelf Sci.*, 56, 1187-1194, 10.1016/S0272-7714(02)00330-x.
- Pastuszak, M., Witek, Z., Nagel, K., Wielgat, M., Grelowski, A., 2005. Role of the Oder estuary (southern Baltic) in transformation of the riverine nutrient loads. *Journal of Marine Systems* 57, 30-54.
- Pastuszak, M., Nagel, K., Grelowski, A., Mohrholz, V. and Zalewski, M., 2003. Nutrient Dynamics in the Pomeranian Bay (Southern Baltic): Impact of the Oder River Outflow. *Estuaries* 26(5): 1238-1254.
- Paerl, H.W., Valdes, L.M., Peierls, B.L., Adolf, J.E., Harding Jr., L.W., 2006. Anthropogenic and climatic influences on the eutrophication of large estuarine ecosystems. *Limnol. Oceanogr.*, 51, 1, part 2, 448-462.
- Paerl, H.W., Huisman, J., 2008. Blooms like it hot. *Science*, 320, 57-58.
- Peierls, B.L., Caraco, N.F., Pace, M.L., Cole, J.J., 1991. Human influence on river nitrogen. *Nature* 350, 386-387.
- Peierls, B.L., Paerl, H.W., 1997. Bioavailability of atmospheric organic nitrogen deposition to coastal phytoplankton. *Limnol. Oceanogr.* 42, 1819-1823.
- Peterson, B. J., and Fry, B., 1987. Stable isotopes in ecosystem studies, *Annual Review of Ecology and Systematics*, 18, 293-320, 10.1146/annurev.ecolsys.18.1.293.
- Petrone, K.C., Richards, J.S., Grierson, P.F., 2009. Bioavailability and composition of dissolved organic carbon and nitrogen in a near coastal catchment of south-western Australia. *Biogeochemistry* 92: 27-40.
- Pettersson, C., Allard, B., Boren, H., 1997. River Discharge of Humic Substances and Humic-bound Metals to the Gulf of Bothnia. *Estuarine, Coastal and Shelf Science* 44: 533-541.
- Philippart, C. J. M., Beukema, J. J., Cadée, G. C., Dekker, R., Goedhart, P. W., van Iperen, J. M.,

- Leopold, M. F., Herman, P. M. J., 2007. Impacts of nutrient reduction on coastal communities. *Ecosystems* 10, 95–118.
- Pilkaityté, R. and Razinkovas, A., 2006. Factors controlling phytoplankton blooms in a temperate estuary: nutrient limitation and physical forcing. *Hydrobiologia* 555: 41-48.
- Pinckney, J.L., Paerl, H.W., Tester, P., and Richardson, T.L., 2001. The Role of Nutrient Loading and Eutrophication in Estuarine Ecology: Environmental Health Perspectives v. 109, p. 699-706.
- Pitkänen, H., Lehtoranta, J., and Peltonen, H., 2008. The Gulf of Finland, in: *Ecology of Baltic Coastal Waters*, edited by: Schiewer, U., Ecological Studies, Springer, Heidelberg.
- Rabalais, N.N., 2002. Nitrogen in Aquatic Ecosystems: *Ambio*, 31, 102-112.
- Rahm, L. and Danielsson, Å., 2007. Spatial heterogeneity of nutrients in the Baltic Proper, Baltic Sea. *Estuarine, Coastal and Shelf Science*, 73(1-2): 268-278.
- Reuter, J. H., 1977. Organic-matter in estuaries, *Chesapeake Science*, 18, 120-121, 10.2307/1350379.
- Risgaard Petersen, N. L. P., Nielsen, S., Rysgaard, D. T., Meyer, R. L., 2003. Application of the isotope pairing technique in sediments where anammox and denitrification coexist. *Limnology and Oceanography: Methods*, 1, 63–73.
- Rolff, C., Elmgren, R., Voss, M., 2008. Deposition of nitrogen and phosphorus on the Baltic Sea: seasonal patterns and nitrogen isotope composition. *Biogeosciences* 5, 1657–1667.
- Rolff, C., and Elmgren, R., 2000. Use of riverine organic matter in plankton food webs of the Baltic Sea, *Mar. Ecol.-Prog. Ser.*, 197, 81-101, 10.3354/meps197081.
- Rummukainen, M., Bergstrom, S., Persson, G., Rodhe, J., Tjernstrom, M., 2004. The Swedish Regional Climate Modelling Programme, SWECLIM: A review. *Ambio* 33, 176-182.
- Rummukainen, M., Räisänen, J., Bjørge, D., Christensen, J.H., Christensen, O.B., Iversen, T., Jylhä, K., Olafson, H., Tuomenvirta, H., 2003. Regional climate scenarios for use in Nordic water resource studies. *Nordic Hydrology* 34, 399-412.
- Santos, I. R., Burnett, W. C., Chanton, J., Mwashote, B., Suryaputra, I. G. N. A., Dittmar, T., 2008. Nutrient biogeochemistry in a gulf of mexico subterranean estuary and groundwater-derived fluxes to the coastal ocean. *Limnology And Oceanography* 53, 2, 705–718.
- Sarmiento, J.L., Gruber, N., 2002. Sinks for anthropogenic carbon. *Physiks Today*, 30-36.
- Savchuk, O.P., 2005. Resolving the Baltic Sea into seven subbasins: N and P budgets for 1991–1999. *Journal of Marine Systems* 56, 1-15.
- Savchuk, O.P., Wulff, F., 2009. Long-term modeling of large-scale nutrient cycles in the entire Baltic Sea. *Hydrobiologia* 629, 209-224.
- Schelske, C.L., Stoermer, E.F., Arbor, A., 1971. Eutrophication, silica depletion, and predicted changes in algal quality in Lake Michigan. *Science* 173, 423 - 424.
- Schelske, C.L., Stoermer, E.F., Conley, D.J., Robbins, J.A., Glover, R.M., 1983. Early eutrophication in the lower Great Lakes: New evidence from biogenic silica in sediments. *Science (Washington)* 222, 320-322.
- Schernewski, G., Neumann, T., 2005. The trophic state of the Baltic Sea a century ago: a model simulation study. *Journal of Marine Systems* 53, 109– 124.
- Schlarbaum, T., Daehnke, K., Emeis, K., 2010. Turnover of combined dissolved organic nitrogen and ammonium in the Elbe estuary/NW Europe: Results of nitrogen isotope investigations. *Marine Chemistry* 119: 91–107.
- Schmidt, G.A., Bigg, G. R., and Rohling, E. J., 1999. Global Seawater Oxygen-18 Database - v1.21: <http://data.giss.nasa.gov/o18data/>.
- Schneider, B., Kaitala, S., Raateoja, M., Sadkowiak, B., 2009. A nitrogen fixation estimate for the Baltic Sea based on continuous pCO<sub>2</sub> measurements on a cargo ship and total nitrogen data. *Continental Shelf Research* 29, 1535-1540.
- Schneider, B., Kuss, J., 2004. Past and present productivity of the Baltic Sea as inferred from pCO<sub>2</sub> data. *Continental Shelf Research* 24: 1611-1622.
- Schuur, E. A. G., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V., Hagemann, S., Kuhry, P., Lafleur, P. M., Lee, H., Mazhitova, G., Nelson, F. E., Rinke, A., Romanovsky, V. E., Shiklomanov, N., Tarnocai, C., Venevsky, S., Vogel, J. G., and Zimov, S. A., 2008. Vulnerability

- of permafrost carbon to climate change: Implications for the global carbon cycle, *Bioscience*, 58, 701-714, 10.1641/b580807.
- Sebilo, M., Billen, G., Mayer, B., Billiou, D., Grably, M., Garnier, J., and Mariotti, A., 2006. Assessing Nitrification and Denitrification in the Seine River and Estuary Using Chemical and Isotopic Techniques: *Ecosystems* 9, 564-577.
- See, J.H., Bronk, D.A., Lewitus, A.J., 2006. Uptake of *Spartina*-derived humic nitrogen by estuarine phytoplankton in nonaxenic and axenic culture. *Limnology and Oceanography* 51: 2290-2299.
- Selman, M., Greenhalgh, S., Diaz, R., Sugg, Z., 2008. Eutrophication and Hypoxia in coastal Areas: A Global Assessment of the state of Knowledge. WRI Policy Note, Water Quality, Eutrophication and Hypoxia, 1.
- Seitzinger, S., Harrison, J.A., Böhlke, J.K., Bouwman, A.F., Lowrance, R., Peterson, B., Tobias, C., Drecht, G.V., 2006. Denitrification across landscapes and waterscapes: A synthesis. *Ecological Applications* 16, 2064–2090 ; DOI: 2010.1890/1051-0761.
- Seitzinger, S., Sanders, R.W., 1999. Atmospheric inputs of dissolved organic nitrogen stimulate estuarine bacteria and phytoplankton. *Limnology Oceanography* 44, 721–730.
- Seitzinger, S.P., Sanders, R.W., 1997. Contribution of dissolved organic nitrogen from rivers to estuarine eutrophication. *Marine Ecology Progress Series* 159, 1-12.
- Seitzinger, S.P., Sanders, R.W., Styles, R., 2002. Bioavailability of DON from natural and anthropogenic sources to estuarine plankton. *Limnology Oceanography* 47, 353–366.
- Seitzinger, S.P., 1988. Denitrification in freshwater and coastal marine ecosystems: Ecological and geochemical significance. *Limnol. Oceanogr.*, 33: 702-724.
- Seitzinger, S.P. et al., 2005. Sources and delivery of carbon, nitrogen, and phosphorus to the coastal zone: An overview of Global Nutrient Export from Watersheds (NEWS) models and their application. *Global Biogeochemical Cycles*, 19(GB4S01): 1-11.
- Seitzinger, S. et al., 2006. Denitrification across landscapes and waterscapes: A synthesis. *Ecol. Appl.* 16, 2064-2090.
- Shaffer, G., Rönner, U., 1984. Denitrification in the Baltic proper deep water. *Deep-Sea Research* 31, 197-220.
- Siegenthaler, U., Sarmiento, J.L., 1993. Atmospheric carbon dioxide and the ocean. *Nature* 365, 119-125.
- Sigleo, A. C., and Macko, S. A., 2002. Carbon and nitrogen isotopes in suspended particles and colloids, Chesapeake and San Francisco estuaries, USA, *Estuar. Coast. Shelf Sci.*, 54, 701-711, 10.1006/ecss.2001.0853.
- Sigman, D.M., Robinson, R., Knapp, A.N., Geen, A.v., McCorkle, D.C., Brandes, J.A., and Thunell, R.C., 2003. Distinguishing between water column and sedimentary denitrification in the Santa Barbara Basin using the stable isotopes of nitrate: *Geochemistry Geophysics Geosystems*, 4, 1-20.
- Sigman, D.M., Altabet, M.A., McCorkle, D.C., Francois, R. and Fischer, G., 1999. The  $\delta^{15}\text{N}$  of nitrate in the southern ocean: Consumption of nitrate in surface waters. *Global Biogeochem. Cycles*, 13(4): 1149-1166.
- Sigman, D.M. et al., 2001. A Bacterial Method for the Nitrogen Isotopic Analysis of Nitrate in Seawater and Freshwater. *Anal. Chem.*, 73: 4145-4153.
- Sigman, D.M. et al., 2005. Coupled nitrogen and oxygen isotope measurements of nitrate along the eastern North Pacific margin. *Global Biogeochemical Cycles*, 19(GB4022): 1-14.
- Sigman, D.M., Altabet, M.A., Michener, R., McCorkle, D.C., Fry, B., Holmes, R.M., 1997. Natural abundance-level measurement of the nitrogen isotopic composition of oceanic nitrate: an adaptation of the ammonia diffusion method. *Marine Chemistry* 57: 227-242.
- Sileika, A.S., Stalnacke, P., Kutra, S., Gaigalis, K. and Berankiene, L., 2006. Temporal and spatial variation of nutrient levels in the Nemunas River (Lithuania and Belarus). *Environmental Monitoring and Assessment* 122: 335-354.

- Silva, S.R., Kendall, C., Wilkison, D.H., Ziegler, A.C., Chang, C.C.Y., and Avanzino, R.J., 2000. A new method for collection of nitrate from fresh water and the analysis of nitrogen and oxygen isotope ratios: *Journal of Hydrology* 228, 22-36.
- Smedberg, E., Humborg, C., Jacobsson, Mörth, C.-M., 2009. Landscape elements and river chemistry as affected by river regulation – a 3-D perspective. *Hydrology and Earth System Sciences* 13, 1597-1606.
- Smedberg, E., Morth, C.M., Swaney, D.P., Humborg, C., 2006. Modeling hydrology and silicon-carbon interactions in taiga and tundra biomes from a landscape perspective: Implications for global warming feedbacks. *Global Biogeochemical Cycles* 20, 15 pp, doi:10.1029/2005GB002567.
- Smith, D.C., Simon, M., Alldredge, A.L., Azam, F., 1992. Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution. *Nature* 359, 139–142.
- Smith, B., Aasa, A., Ahas, R., Blenckner, T., Callaghan, T., de Chazal, J., Humborg, C., Jönsson, A.M., Kellomäki, S., Kull, A., Lehikoinen, E., Mander, Ü., Nöges, P., Nöges, T., Rounsevell, M., Sofiev, M., Tryjanowski, P., Wolf, A., , 2008. Climate-related change in terrestrial and freshwater ecosystems, In: (Eds.), BACC Author Group, Assessment of Climate Change for the Baltic Sea Basin. Springer-Verlag, Germany, 474 pp.
- Soetaert, K., Middelburg, J.J., Heip, C., Meire, P., Damme, S.V., Maris, T., 2006. Long-term change in dissolved inorganic nutrients in the heterotrophic Scheldt estuary (Belgium, The Netherlands). *Limnol. Oceanogr.* 51, 409–423.
- Sommer, F., Hansen, T., Sommer, U., 2006. Transfer of diazotrophic nitrogen to mesozooplankton in Kiel Fjord, Western Baltic Sea: a mesocosm study. *Marine Ecology Progress Series* 324: 105–112.
- Spokes, L., Jickells, T., Weston, K., Gustafsson, B.G., Johnsson, M., Liljebladh, B., Conley, D., Ambelas-Skjødth, C., Brandt, J., Carstensen, J., Christiansen, T., Frohn, L., Geernaert, G., Hertel, O., Jensen, B., Lundsgaard, C., Markager, S., W. Martinsen, Møller, B., Pedersen, B., Sauerberg, K., Sørensen, L.L., Hasager, C.C., Sempreviva, A.M., Pryor, S.C., Lund, S.W., Larsen, S., Tjernström, M., Svensson, G., Zagar, M., 2006. MEAD: An interdisciplinary study of the marine effects of atmospheric deposition in the Kattegat. *Environmental Pollution* 140, doi:10.1016/j.envpol.2005.1008.1010, 1453-1462.
- Stålnacke, P., Grimvall, A., Sundblad, K. and Tonderski, A., 1999. Estimation of riverine loads of nitrogen and phosphorus to the Baltic Sea, 1970-1993. *Environmental Monitoring and Assessment*, 58(2): 173-200.
- Stedmon, C.A., S. Markager, M. Sondergaard, T. Vang, A. Laubel, N. H. Borch, and A. Windelin, 2006. Dissolved Organic Matter (DOM) Export to a Temperate Estuary: Seasonal Variations and Implications of Land Use. *Estuaries and Coasts* 29, 388–400.
- Steinberg, D.K., Carlson, C.A., Bates, N.R. , Goldthwait, S.A., Madin, L.P., Michaels, A.F., 2000. Zooplankton vertical migrations and the active transport of dissolved organic and inorganic carbon in the Sargasso Sea. *Deep-Sea Research I* 47, 137-158.
- Stepanaukas, R., Jorgensen, N.O.G., Eigaard, O.R., Zvikas, A., Tranvik, L.J., Leonardson, L., 2002. Summer inputs of riverine nutrients to the Baltic Sea: Bioavailability and eutrophication relevance. *Ecological Monographs* 72, 579-597
- Stepanaukas, R., Laudon, H., Jørgensen, N.O.G., 2000. High DON bioavailability in boreal streams during a spring flood. *Limnology Oceanography* 45, 1298–1307.
- Stepanaukas, R., Leonardson, L., Tranvik, L.J., 1999. Bioavailability of wetland-derived DON to freshwater and marine bacterioplankton. *Limnology Oceanography* 44, 1477–1485.
- Stepanaukas, R., Edling, H., Tranvik, L.J., 1999b. Differential dissolved organic nitrogen availability and bacterial aminopeptidase activity in limnic and marine waters. *Microbial Ecology* 38: 264-272.
- Stepanaukas, R., Moran, M. A., Bergamasch, B. A., and Hollibaugh, J. T., 2005. Sources, bioavailability, and photoreactivity of dissolved organic carbon in the Sacramento-San Joaquin River Delta, *Biogeochemistry*, 74, 131-149, 10.1007/s10533-004-3361-2.

- Stigebrandt, A., 2001. Physical Oceanography of the Baltic Sea, In: Wulff, F.V., Rahm, L.A.Larsson, P. (Eds.), *A Systems Analysis of the Baltic Sea*. Springer-Verlag, Berlin Heidelberg, pp. 19-74.
- Striegl, R.G., Dornblaser, M.M., Aiken, G.R., Wickland, K.P., Raymond, P.A., 2007. Carbon export and cycling by the Yukon, Tanana, and Porcupine rivers, Alaska, 2001-2005. *Water Resources Research* 43.
- Struck, U., Emeis, K.-C., Voss, M., Christiansen, C. and Kunzendorf, H., 2000. Records of southern and central Baltic Sea eutrophication in  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  of sedimentary organic matter. *Marine Geology* 164: 157-171.
- Struck, U., Pollehne, F., Bauerfeind, E. and v. Bodungen, B., 2004. Sources of nitrogen for the vertical particle flux in the Gotland Sea (Baltic Proper)- Results from sediment trap studies. *Journal of Marine Systems*, 45(1-2): 91-101.
- Sutka, R.L., Ostrom, N.E., Ostrom, P.H., and Phanikumar, M.S., 2004. Stable nitrogen isotope dynamics of dissolved nitrate in a transect from the north Pacific subtropical gyre to the eastern tropical north Pacific: *Geochimica et Cosmochimica Acta*, 69, 517-527.
- Suttle, C.A., 1994. The significance of viruses to mortality in aquatic microbial communities. *Microbial Ecology* 28, 237-243.
- Swanson, K.L., Tsonis, A.A., 2009. Has the climate recently shifted? *Geophysical Research Letters* 36, L06711, doi:10.1029/2008GL037022.
- Sweitzer, J., Langaas, S., Folke, C., 1996. Land use and population density in the Baltic Sea Drainage Basin: A GIS database. *Ambio* 25: 191-98.
- Syvitski, J.P.M., Vörösmarty, C.J., Kettner, A.J., and Green, P., 2005. Impact of Humans on the Flux of Terrestrial Sediment to the Global Coastal Ocean: *Science* 308.
- Tamminen, T., Andersen, T., 2007. Seasonal phytoplankton nutrient limitation patterns as revealed by bioassays over Baltic Sea gradients of salinity and eutrophication. *Marine Ecology Progress Series* 340, 121-138.
- Tamminen, T., Irmisch, A., 1996. Urea uptake kinetics of a midsummer planktonic community on the SW coast of Finland. *Marine Ecology Progress Series* 130: 201-211.
- Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S., 2009. Soil organic carbon pools in the northern circumpolar permafrost region, *Glob. Biogeochem. Cycle*, 23, 11, Gb2023 10.1029/2008gb003327.
- Thibodeaux, L.J. and Boyle, J.D., 1987. Bedform-generated convective transport in bottom sediment. *nature*, 325(6102): 341-343.
- Thomas, H., Ittekkot, V., Osterroht, C., Schneider, B., 1999. Preferential recycling of nutrients - the ocean's way to increase new production and to pass nutrient limitation? *Limnology And Oceanography* 44, 8, 1999-2004.
- Thomson, D.W.J., Wallace, J.M., 1998. The Arctic Oscillation signature in the wintertime geopotential height and temperature field. *Geophysical Research Letters* 25, 1297-1300.
- Timperley, M.H., R. J. Vigor-Brown, M. Kawashima, and M. Ishigami, 1985. Organic nitrogen compounds in atmospheric precipitation: Their chemistry and availability to phytoplankton. *Can. J. Fish. Aquat. Sci.* 42, 1171-1177.
- Tobias, C.R., Harvey, J.W., and Anderson, I.C., 2001. Quantifying groundwater discharge through fringing wetlands to estuaries: Seasonal variability, methods comparison, and implications for wetland-estuary exchange. *Limnol. Oceanogr.* 46, 604-615.
- Tranvik, L.J., Jansson, M., 2002. Climate change - Terrestrial export of organic carbon. *Nature* 415, 861-862.
- Vahtera, E., Conley, D.J., Gustafsson, B.G., Kuosa, H., Pitkänen, H., Savchuk, O.P., Tamminen, T., Viitasalo, M., Voss, M., Wasmund, N., Wulff, F., 2007. Internal Ecosystem Feedbacks Enhance Nitrogen-fixing Cyanobacteria Blooms and Complicate Management in the Baltic Sea. *Ambio* 36, 186-194.
- Valiela, I., Costa, J. Foreman, K., Teal, J.M., Howes, B., and Aubrey, D., 1990. Transport of groundwater-borne nutrients from watersheds and their effects on coastal waters. *Biogeochemistry* 10, 177-197.

- Van Engeland, T.V., Soetaert, K., Knuijt, A., Laane, R.W.P.M., Middelburg, J.J., 2010. Dissolved organic nitrogen dynamics in the North Sea: A time series analysis (1995-2005). *Estuarine, Coastal and Shelf Science* 89: 31-42.
- Van den Meersche, K., Van Rijswijk, P., Soetaert, K., and Middelburg, J. J., 2009. Autochthonous and allochthonous contributions to mesozooplankton diet in a tidal river and estuary: Integrating carbon isotope and fatty acid constraints, *Limnol. Oceanogr.*, 54, 62-74, 10.4319/lo.2009.54.1.0062.
- Veuger, B., Middelburg, J.J., Boschker, H.T.S., Nieuwenhuize, J., Rijswijk, P.v., Rochelle-Newall, E.J., Navarro, N., 2004. Microbial uptake of dissolved organic and inorganic nitrogen in Randers Fjord. *Estuarine, Coastal and Shelf Science* 61, 507–515.
- Veuger, B., Middelburg, J. J., 2007. Incorporation of nitrogen from amino acids and urea by benthic microbes: role of bacteria versus algae and coupled incorporation of carbon. *Aquatic Microbial Ecology* 48, 1, 35–46.
- Vitousek, P.M. et al., 1997. Human alteration of the global nitrogen cycle: Sources and consequences. *Ecological Applications*, 7(3): 737-750.
- Voss, M., J. W. Dippner, et al., 2001. Nitrogen isotope patterns in the oxygen-deficient waters of the Eastern Tropical North Pacific Ocean. *Deep-Sea Research Part I-Oceanographic Research Papers* 48(8): 1905-1921.
- Voss, M., M. A. Altabet, et al., 1996.  $\delta$  N-15 in sedimenting particles as indicator of euphotic-zone processes. *Deep-Sea Research Part I-Oceanographic Research Papers* 43(1): 33-47.
- Voss, M., Deutsch, B., Elmgren, R., Humborg, C., Kuuppo, P., Pastuszak, M., Rolff, C., Schulte, U., 2006. Source identification of nitrate by means of isotopic tracers in the Baltic Sea catchments. *Biogeosciences* 3, 663-676.
- Voss, M., Deutsch, B., Liskow, I., Pastuszak, M., Schulte, U., Sitek, S., 2010. Nitrogen retention in the Szczecin Lagoon, Baltic Sea. *Isotopes in Environmental and Health Studies* 47, 1–15, DOI: 10.1080/10256016.10252010.10503895.
- Voss, M., Emeis, K.-C., Hille, S., Neumann, T., Dippner, J.W., 2005a. Nitrogen cycle of the Baltic Sea from an isotopic perspective. *Global Biogeochemical Cycles* 19, 1-16, doi:10.1029/2004GB002338.
- Voss, M., Larsen, B., Leivuori, M., Vallius, H., 2000. Stable isotope signals of eutrophication in coastal Baltic Sea sediments. *Journal of Marine Systems (Special Issue)* 25, 287-298.
- Voss, M., Liskow, I., Pastuszak, M., Ruess, D., Dippner, J.W., 2005b. Riverine Discharge into a Coastal Bay: A stable Isotope study in the Gulf of Gdańsk, Baltic Sea. *Journal of Marine Systems* 57, 127-145.
- Voss, M., Dippner, J. W., Humborg, C., Hurdler, J., Korth, F., Neumann, T., Schernewski, G., and Venohr, M., 2011. History and scenarios of future development of Baltic Sea eutrophication. *Estuarine, Coastal and Shelf Science*, 92(3): 307-322.
- Voss, M. and Struck, U., 1997. Stable nitrogen and carbon isotopes as indicator of eutrophication of the Oder river (Baltic Sea) *Marine Chemistry* 59: 35-49.
- Walvoord, M.A., Striegl, R.G., 2007. Increased groundwater to stream discharge from permafrost thawing in the Yukon River basin: Potential impacts on lateral export of carbon and nitrogen. *Geophysical Research Letters* 34, L12402, doi:10.1029/2007GL030216.
- Wankel, S.D., Kendall, C., Francis, C.A. and Paytan, A., 2006. Nitrogen sources and cycling in the San Francisco Bay Estuary: A nitrate dual isotopic composition approach. *Limnol. Oceanogr.*, 51(4): 1654-1664.
- Wankel, S.D., Kendall, C., Pennington, J.T., Chavez, F.P. and Paytan, A., 2007. Nitrification in the euphotic zone as evidenced by nitrate dual isotopic composition: Observations from Monterey Bay, California. *Global Biogeochemical Cycles*, 21: 1-13.
- Wannicke, N., Koch, B.P., Voss, M., 2009. Release of fixed N<sub>2</sub> and C as dissolved compounds by *Trichodesmium erythreum* and *Nodularia spumigena* under the influence of high light and high nutrient (P). *Aquatic Microbiological Ecology* 57: 175-189.

- Wasmund, N., Uhlig, S., 2003. Phytoplankton trends in the Baltic Sea. *ICES Journal of Marine Science* 60, 177-186.
- Wasmund, N., Voss, M., Lochte, K., 2001. Annual nitrogen fixation patterns in the Baltic Proper. *Marine Ecology Progress Series* 214, 1-14.
- Wasmund, N., Nausch, G., Schneider, B., Nagel, K., Voss, M., 2005. Comparison of nitrogen fixation and primary production rates by different methods: A study in the Baltic Proper. *Marine Ecology Progress Series* 297: 23-31.
- Wiegner, T.N., Seitzinger, S.P., Glibert, P.M., Bronk, D.A., Bronk, D.A., 2006. Bioavailability of dissolved organic nitrogen and carbon from nine rivers in the eastern United States. *Aquatic Microbial Ecology* 43, 277–287.
- Wiegner, T. N., and Seitzinger, S. P., 2004. Seasonal bioavailability of dissolved organic carbon and nitrogen from pristine and polluted freshwater wetlands, *Limnol. Oceanogr.*, 49, 1703-1712.
- Wiegner, T. N., Tubal, R. L., MacKenzie, R. A., 2009. Bioavailability and export of dissolved organicmatter from a tropical river during base- and stormflow conditions. *Limnology and Oceanography* 54, 4, 1233–1242.
- Williams, C. J., Yamashita, Y., Wilson, H. F., Jaffe, R., and Xenopoulos, M. A., 2010. Unraveling the role of land use and microbial activity in shaping dissolved organic matter characteristics in stream ecosystems, *Limnol. Oceanogr.*, 55, 1159-1171, 10.4319/lo.2010.55.3.1159.
- Winsor, P., Rodhe, J., Omstedt, A., 2001. Baltic Sea ocean climate: an analysis of 100 yr of hydrographic data with focus on the freshwater budget. *Climate Research* 18, 5-15.
- Witek, Z., Drgas, A., Ameryk, A. and Ochocki, S., 2001. Production and mineralization of organic matter in the Pomeranian Bay. *Bulletin of the sea fisheries institute*, 3(154): 49-69.
- Wu, J., Calvert, S.E. and Wong, C.S., 1997. Nitrogen isotope variations in the subarctic northeast Pacific: relationships to nitrate utilization and trophic structure. *Deep Sea Research Part I: Oceanographic Research Papers*, 44(2): 287-314.
- Wulff, F., Humborg, C., Medina, M.R., Mörtz, C.-M., Savchuk, O., and Sokolov, A., 2009. Revision of the country allocation of nutrient reductions in the Baltic Sea Action Plan, *Baltic Nest Institute, Stockholm Resilience Center: Stockholm, Baltic Nest Institute*, 1-47.
- Zehr, J. P. , Kudela, R. M., 2011. Nitrogen Cycle of the Open Ocean: From Genes to Ecosystems. *Annu. Rev. Mar. Sci.* 3, 197–225.
- Zillén, L., Conley, D.J., Andrén, T., Andrén, E., Björck, S., 2008. Past occurrences of hypoxia in the Baltic Sea and the role of climate variability, environmental change and human impact. *Earth-Science Reviews* 91, 77-92.
- Zimov, N. S., Zimov, S. A., Zimova, A. E., Zimova, G. M., Chuprynin, V. I., and Chapin, F. S., 2009. Carbon storage in permafrost and soils of the mammoth tundra-steppe biome: Role in the global carbon budget, *Geophys. Res. Lett.*, 36, L02502 10.1029/2008gl036332.
- Zimov, S. A., Schuur, E. A. G., and Chapin, F. S., 2006. Permafrost and the global carbon budget, *Science*, 312, 1612-1613, 10.1126/science.1128908.