

Documentation

Dataset for Manuscript "High-Order Harmonic Generation in an Organic Molecular Crystal"

Falk-Erik Wiechmann, Samuel Schöpa, Lina Bielke, Svenja Rindelhardt,
Serguei Patchkovskii, Felipe Morales, Maria Richter,
Dieter Bauer, Franziska Fennel

1 General Information

Dataset title: Dataset for Manuscript "High-Order Harmonic Generation in an Organic Molecular Crystal"

Authors/Creators: Falk-Erik Wiechmann^{1,2}, Samuel Schöpa¹, Lina Bielke¹, Svenja Rindelhardt^{1,2}, Serguei Patchkovskii³, Felipe Morales³, Maria Richter³, Dieter Bauer^{1,2}, Franziska Fennel^{1,2}

Affiliations: ¹Institute of Physics, University of Rostock, Albert-Einstein-Str. 23, 18059 Rostock, Germany
²Department of Life, Light & Matter, University of Rostock, 18051 Rostock, Germany
³Max Born Institute, Max-Born-Str. 2A, 12489 Berlin, Germany

E-Mails: dieter.bauer@uni-rostock.de
franziska.fennel@uni-rostock.de

Date: October 2025

Type: Measurements, Simulations, Analysis Results

Language: English

Rights: CC BY-NC 4.0

DOI: https://doi.org/10.18453/rosdok_id00004817

Funding: We acknowledge funding from the Deutsche Forschungsgemeinschaft via SFB 1477 'Light-Matter Interactions at Interfaces' (project no. 441234705).

2 Description

This document provides the data, which is discussed in the publication [1].

3 File Format

The data is provided in the form of ".txt"-files containing the research data in tables. It can be easily accessed in python with the numpy library using the "numpy.loadtxt("filename.txt")" command or in matlab with "readtable("filename.txt")". All entries use double-precision floats and row-major ordering.

4 Repository Structure

The data is grouped into the following sub-folders and files:

```
HHG_OMC_Repository
├── Absorption_Spectra
│   ├── OD.txt
│   ├── Polarization.txt
│   └── Wavelength.txt
├── Polarization_Dependent_Yields
│   ├── Yield_Experiment.txt
│   ├── Yield_MR_CIS.txt
│   ├── Yield_TB.txt
│   ├── Yield_TB_flipped_t2_0_01.txt
│   └── Yield_TB_flipped_t2_0_0001.txt
├── Tight_Binding_Flipping_Analysis
│   ├── phi_of_maximum_yield.txt
│   └── t2_values.txt
├── Single_HHG_Spectra
│   ├── Spectrum_Experiment.txt
│   ├── Spectrum_MR_CIS.txt
│   └── Spectrum_TD_DFT.txt
└── Documentation.pdf
```

4.1 Absorption_Spectra

Includes the data supporting Extended Data Figure 1.

OD.txt	Optical density of the measured polarizations and wavelengths
Polarization.txt	Polarization of the driving field in [°]
Wavelength.txt	Wavelength in [nm]

4.2 Polarization_Dependent_Yields

Includes the data supporting all shown polar plots, i.e. Fig. 2, 3 and Extended Data Figure 3. In all data files we provide the harmonic yield and the corresponding laser polarization.

Yield_Experiment.txt	The first row contains the laser polarization, and the following rows contain the experimentally measured harmonic yield of harmonic order 3 to 15 (Fig. 2, 3)
Yield_MR_CIS.txt	The first row contains the laser polarization, and the following rows contain the harmonic yields of order 1 to 13 for the combined emission of two uncoupled molecules calculated with MR-CIS (Fig. 3)
Yield_TD_DFT.txt	The first row contains the laser polarization, and the following rows contain the harmonic yield of harmonic order 1 to 13 obtained with the TD-DFT calculation for the pentacene crystal (Fig. 3)
Yield_TB.txt	The first row contains the laser polarization, and the following rows contain the harmonic yield of harmonic order 1 to 13 obtained with the 2D-tight-binding model calculation (Fig. 3)
Yield_TB_flipped_t2_0_01.txt	The first row contains the laser polarization, and the following rows contain the harmonic yield of harmonic order 1 to 13 obtained with the 2D-tight-binding model calculation, that has been rotated, for a coupling $t_2 = 0.01$ (Extended Data Figure 3)
Yield_TB_flipped_t2_0_0001.txt	The first row contains the laser polarization, and the following rows contain the harmonic yield of harmonic order 1 to 13 obtained with the 2D-tight-binding model calculation, that has been rotated, for a coupling $t_2 = 0.0001$ (Extended Data Figure 3)

4.3 Tight_Binding_Flipping_Analysis

Includes the data supporting the plots showing the t_2 dependent polarization dependence of the modified tight-binding model, i.e. Fig. 4 and Extended Data Figure 3.

phi_of_maximum_yield.txt	Laser polarization φ for which the harmonic yield is maximized as a function of t_2 for harmonic 1 - 13
t2_values.txt	Values of the tight-binding parameter t_2

4.4 Single_HHG_Spectra

Includes the data supporting the experimental HHG spectra shown in Fig.1 and the comparison of the individual HHG spectra shown in Extended Data Figure 2.

Spectrum_Experiment.txt	The first row contains the energy axis in units of ω/ω_0 , and the second row contains the experimentally measured harmonic spectra. Note, that the spectrum around the third harmonic was measured with a different spectrometer and has been shifted by a factor of 0.3. The electric field is polarized at an angle $\varphi = XXX^\circ$, corresponding a polarization parallel to the direction of a nearest neighbor.
Spectrum_MR_CIS.txt	The first row contains the energy axis in units of ω/ω_0 , and the second row contains the harmonic yield obtained in the MR-CIS simulation for the individual molecules. The electric field is polarized at an angle $\varphi = 130^\circ$, corresponding to a polarization parallel to the long molecular axis.
Spectrum_TD_DFT.txt	The first row contains the energy axis in units of ω/ω_0 , and the second row contains the harmonic yield obtained in the TD-DFT simulation for the crystal. The electric field is polarized at an angle $\varphi = 129.375^\circ$, corresponding a polarization parallel to the direction of a nearest neighbor.

References

- [1] Falk-Erik Wiechmann, Samuel Schöpa, Lina Bielke, Svenja Rindelhardt, Serguei Patchkovskii, Felipe Morales, Maria Richter, Dieter Bauer, Franziska Fennel, *High-Order Harmonic Generation in an Organic Molecular Crystal*, Nature Communications (submitted)